

MEASURABLE SCHUR MULTIPLIERS AND COMPLETELY BOUNDED MULTIPLIERS OF THE FOURIER ALGEBRAS

NICO SPRONK

ABSTRACT. Let G be a locally compact group, $L^p(G)$ be the usual L^p -space for $1 \leq p \leq \infty$ and $A(G)$ be the Fourier algebra of G . Our goal is to study, in a new abstract context, the completely bounded multipliers of $A(G)$, which we denote $M_{cb}A(G)$. We show that $M_{cb}A(G)$ can be characterised as the “invariant part” of the space of (completely) bounded normal $L^\infty(G)$ -bimodule maps on $\mathcal{B}(L^2(G))$, the space of bounded operator on $L^2(G)$. In doing this we develop a function theoretic description of the normal $L^\infty(X, \mu)$ -bimodule maps on $\mathcal{B}(L^2(X, \mu))$, which we denote by $V^\infty(X, \mu)$, and name the *measurable Schur multipliers* of (X, μ) . Our approach leads to many new results, some of which generalise results hitherto known only for certain classes of groups. Those results which we develop here are a uniform approach to obtaining the functorial properties of $M_{cb}A(G)$, and a concrete description of a standard predual of $M_{cb}A(G)$.

1. INTRODUCTION

Let G be a locally compact group, and $A(G)$ the Fourier algebra of G from [17].

In [25], Herz defined the space of *Herz-Schur multipliers*, $B_2(G)$. He claimed this space to coincide with the space he denoted $\mathcal{V}(G)$ in his article [24], but offered no proof of this fact. The *completely bounded multipliers* of the Fourier algebra, $M_{cb}A(G)$, were defined by De Canneire and Haagerup in [8]. It was shown by Božeko and Fendler, [7], that $M_{cb}A(G) = B_2(G)$, isometrically. Unfortunately, this result relies on unpublished work of Gilbert [20]. However, there is a proof of the fact that $M_{cb}A(G) = B_2(G)$ due to Jolissaint [28], which relies on the representation theorem for completely bounded maps applied to the reduced C^* -algebra, $C_r^*(G)$.

Date: February 1, 2008.

2000 *Mathematics Subject Classification.* Primary 46L07, 43A30, 43A15, 46A32, 22D10, 22D12; Secondary 22D25, 43A20, 43A07. *Key words and phrases.* Fourier algebra, operator space, Haagerup tensor product, completely bounded multipliers, Schur multipliers, similarity degree.

This work was supported at various stages by an NSERC PGS B and by Ontario Graduate Scholarships.

In this paper we prove an analogue of the fact that $M_{cb}A(G) = B_2(G)$. In fact, our result obtains that fact in the case that G is a discrete group, and improves upon it in the respect that we obtain a complete isometry, where both $M_{cb}A(G)$ and $B_2(G)$ are given natural operator space structures. Moreover, we show that the natural module action of $M_{cb}A(G)$ on the von Neumann algebra generated by the left regular representation, $VN(G)$, extends completely isometrically to a normal action of $M_{cb}A(G)$ on all of $\mathcal{B}(L^2(G))$. In fact, the image of $M_{cb}A(G)$ in the (completely) bounded normal maps on $\mathcal{B}(L^2(G))$ consists of $L^\infty(G)$ -bimodule maps, and can be characterised as a certain intrinsic “invariant part” amongst those maps. We feel that our approach is interesting in and of itself since it makes systematic use of modern operator space theoretic techniques, pioneered by Blecher, Effros, Haagerup, Paulsen, Ruan and Smith, amongst others. It thus gives a starting point for obtaining representations of completely multipliers of Kac algebras, after [30] or of multipliers of locally compact quantum groups after Vaes et al (see [49]).

To support our approach, in Section 3 we describe the space of *measurable Schur multipliers*, $V^\infty(X, \mu)$, for any suitable measure space (X, μ) . We develop the theory of $V^\infty(X, \mu)$ fully and systematically. Our main tools for doing this are the results of Smith [43] and Blecher and Smith [5] which characterise the normal completely bounded maps on the space of all bounded operators on a Hilbert space, which are module maps over a von Neumann algebra. We make great use of the weak* Haagerup tensor product of two von Neumann algebras, which is described in [5]. In particular, we note that $V^\infty(X, \mu)$ obtains a natural operator space structure. Subordinate to the theory of measurable Schur multipliers, we develop a theory of “continuous” Schur multipliers, which is based on the theory of the extended Haagerup tensor product of Effros and Ruan in [14] (see also [13]).

Letting m now denote the left Haar measure on our locally compact group G , we let $V^\infty(G) = V^\infty(G, m)$. We define the “invariant part”, $V_{\text{inv}}^\infty(G)$, of $V^\infty(G)$. Our space $V_{\text{inv}}^\infty(G)$ is essentially the space $\mathcal{V}(G)$ of [24]. Our main result is that $V_{\text{inv}}^\infty(G) \cong M_{cb}A(G)$ completely isometrically. Thus we obtain a natural extension of the module action of $M_{cb}A(G)$ on $VN(G)$ to $\mathcal{B}(L^2(G))$. This is the content of Section 5. (It should be noted that the isometric identification $V_{\text{inv}}^\infty(G) \cong M_{cb}A(G)$ has been previously known to Haagerup [21]. This fact was discovered by the author after the present article was completed. However, it is not clear how the methods of that article can be adapted to give a complete isometry.)

As an immediate application of our results, we have a systematic approach to determining the functorial properties of $M_{cb}A(G)$. In addition, we can identify some circumstances in which a subspace of the Fourier-Stieltjes algebra, $B(G)$, acts completely isometrically as multipliers of $A(G)$. Another application we obtain is a concrete description of the predual of $M_{cb}A(G)$, $Q(G)$, whose existence is recognised in [8]. In the case where G is discrete, Pisier [40] implicitly uses the structure of $Q(G)$ in his partial solution to

the similarity problem for representations of groups. These applications are studied in Section 6.

We note that there are further applications of our results and methods which are not explored in this note. As an example, the author and L. Turowska, in [45], use some of the methods of the current note to show that if G is compact, then $A(G)$ naturally imbeds into $\mathcal{C}(G) \otimes^\gamma \mathcal{C}(G)$ (projective tensor product of the continuous functions on G with itself). This is used to generalise, to arbitrary compact groups, results on parallel spectral synthesis due to Varopoulos [50] for compact Abelian groups; and to generalise some results on operator synthesis due to Froelich [18].

We should also point out that if G is Abelian, there are related results to ours by Størmer [47]. In that article it is shown that there is an isometric representation of the measure algebra, $M(G)$, in $\mathcal{B}^\sigma(L^2(G))$, which extends convolution on $L^\infty(G)$. This is generalised to arbitrary locally compact groups by Ghahramani [19]. Neufang [34] has shown that this isometric representation has its range in $\mathcal{CB}^\sigma(L^2(G))$. Some of the connections between these results and those of the present article are being explored by Neufang and the author [35].

Sections 2 and 4 are included to support the exposition of this article, but include results of independent interest. In Section 2, a uniform approach is developed for defining the weak* and extended Haagerup tensor products. In Section 4 some basic results about completely bounded multipliers are reviewed, and a theorem of Walter [52] is improved.

This article represents a significant portion of the author's doctoral thesis [44]. The author would like to express his gratitude to his advisor, Brian Forrest, for support and guidance throughout his tenure as a graduate student at the University of Waterloo. His gratitude extends to other members of the faculty there, including Ken Davidson, Kathryn Hare and Andu Nica. The author would also like to acknowledge his fellow graduate student Peter Wood, and the excellent staff in the Pure Math department.

1.1. Notation. If $\mathcal{X}$ is a complex normed vector space, we will let $b_1(\mathcal{X})$ denote the closed unit ball, $\mathcal{X}^*$ denote the space of continuous linear functionals on $\mathcal{X}$, i.e. the dual space, and $\mathcal{B}(\mathcal{X})$ denote the normed algebra of bounded linear operators on $\mathcal{X}$. If $T \in \mathcal{B}(\mathcal{X})$, we let $T^* : \mathcal{X}^* \rightarrow \mathcal{X}^*$ denote the adjoint operator. If $\mathcal{H}$ is a Hilbert space and $x \in \mathcal{B}(\mathcal{H})$, we let $x^* : \mathcal{H} \rightarrow \mathcal{H}$ denote its Hilbertian adjoint.

If $\mathcal{X}$ is a dual space, with established predual $\mathcal{X}_*$, we let $\mathcal{B}^\sigma(\mathcal{X})$ denote the space of linear operators on $\mathcal{X}$ which are continuous in the $\sigma(\mathcal{X}, \mathcal{X}_*)$ topology. Hence if $\mathcal{M}$ is a von Neumann algebra, $\mathcal{B}^\sigma(\mathcal{M})$ denotes the algebra of normal operators on $\mathcal{M}$.

If X is a locally compact Hausdorff space, let $\mathcal{C}_b(X)$ denote the C^* -algebra of continuous complex-valued bounded functions on X , with norm $\|\varphi\|_\infty = \sup\{|\varphi(x)| : x \in X\}$. We will let $\mathcal{C}_0(X)$ denote the norm closure of the space of compactly supported continuous complex-valued functions on X .

Our standard reference for operator spaces is [16]. However we were also influenced by the points of view in [3] and [4]. There is a good summary of operator spaces in [41].

Given an operator space $\mathcal{V}$, we let $M_n(\mathcal{V})$ denote the space of $n \times n$ matrices with entries in $\mathcal{V}$. We also let $\mathcal{CB}(\mathcal{V})$ denote the algebra of completely bounded linear maps on $\mathcal{V}$, i.e. maps $T : \mathcal{V} \rightarrow \mathcal{V}$ for which the amplifications, $T^{(n)} : M_n(\mathcal{V}) \rightarrow M_n(\mathcal{V})$, satisfy $\|T\|_{cb} = \sup\{\|T^{(n)}\| : n \in \mathbb{N}\} < +\infty$. We will assign $\mathcal{CB}(\mathcal{V})$ and $\mathcal{V}^*$ the operator space from [3] or [16, Sec. 3.2].

If $\mathcal{X}$ is a normed vector space, we let $\max \mathcal{X}$ denote the maximal operator space with underlying normed space $\mathcal{X}$, and we denote by $\min \mathcal{X}$ the minimal one.

We will make substantial use of tensor products here. If $\mathcal{V}$ and $\mathcal{W}$ are complete operator spaces, let $\mathcal{V} \hat{\otimes} \mathcal{W}$ denote their *operator projective tensor product*, and $\mathcal{V} \otimes^h \mathcal{W}$ their *Haagerup tensor product*. We will often write $\mathcal{V} \otimes_{\wedge} \mathcal{W}$, respectively $\mathcal{V} \otimes_h \mathcal{W}$, for the algebraic tensor product $\mathcal{V} \otimes \mathcal{W}$, but with the operator space structure given by the operator projective tensor norms, respectively Haagerup, tensor norms. For Banach spaces $\mathcal{X}$ and $\mathcal{Y}$, $\mathcal{X} \otimes^{\gamma} \mathcal{Y}$ will denote their (Banach space) *projective tensor product*. If $\mathcal{M}$ and $\mathcal{N}$ are von Neumann algebras we will denote the *von Neumann tensor product* by $\mathcal{M} \overline{\otimes} \mathcal{N}$.

2. THE WEAK* AND EXTENDED HAAGERUP TENSOR PRODUCTS

In this article we will make extensive use of the weak* Haagerup tensor product of [5] and the extended Haagerup tensor product of [14] (see also [13]). However, for our applications we will not want to assume that we can work with operator spaces over a separable Hilbert space. Moreover, we want a representation of the extended Haagerup tensor product modeled after one of the weak* Haagerup tensor product in [5, Theo. 3.1]. Hence we will briskly develop a theory of these tensor products in a form that fits our needs.

First we need a description of infinite matrices over an operator space. Let $\mathcal{V}$ be an operator space and I_0 and J_0 be finite index sets. The set of $I_0 \times J_0$ -matrices, $M_{I_0, J_0}(\mathcal{V})$, can be normed in an unambiguous way with no regard to having an ordering on I_0 or J_0 ; see [16, (2.1.5)]. If I and J are infinite index sets, let $M_{I, J}(\mathcal{V})$ denote the set of $I \times J$ matrices with entries in $\mathcal{V}$. If I_0 and J_0 are finite subsets of I and J , and $v \in M_{I, J}(\mathcal{V})$, let v_{I_0, J_0} be the matrix in $M_{I_0, J_0}(\mathcal{V})$ given by restricting to the indices from I_0 and J_0 . We let

$$M_{I, J}(\mathcal{V}) = \left\{ v \in M_{I, J}(\mathcal{V}) : \|v\| = \sup_{\substack{I_0 \subset I, J_0 \subset J \\ \text{finite}}} \|v_{I_0, J_0}\| < +\infty \right\}.$$

As in [16, Sec. 10.1], this is an operator space. For example, if $\mathcal{H}$ is a Hilbert space, then $M_{I,J}(\mathcal{B}(\mathcal{H})) \cong \mathcal{B}(\mathcal{H}^J, \mathcal{H}^I)$, in the usual way. In particular, $M_{I,J} = M_{I,J}(\mathbb{C}) \cong \mathcal{B}(\ell^2(J), \ell^2(I))$.

If $\mathcal{W}$ is another operator space and $S \in \mathcal{CB}(\mathcal{V}, \mathcal{W})$, then the amplification $S^{(I,J)} : M_{I,J}(\mathcal{V}) \rightarrow M_{I,J}(\mathcal{W})$ can be easily seen to have its range in $M_{I,J}(\mathcal{W})$ and to be completely bounded with $\|S^{(I,J)}\| = \|S^{(I,J)}\|_{cb} = \|S\|_{cb}$. Moreover, $S^{(I,J)}$ is a complete isometry if S is. However, if S is a complete quotient map, $S^{(I,J)}$ may not be a (complete) quotient map; it is such, though, if $\mathcal{V}$ and $\mathcal{W}$ are operator dual spaces and S is the adjoint of a complete isometry between their preduals.

Usually, a matrix v in $M_{I,J}(\mathcal{V})$ will be written $v = [v_{ij}]$. However, for row matrices v in $M_{1,J}(\mathcal{V})$ and column matrices w in $M_{I,1}(\mathcal{V})$ we will often write $v = [v_j]$ and $w = [w_i]$, i.e. $v_j = v_{1j}$ and $w_i = w_{i1}$ for each i and j .

2.1. The Fubini-Haagerup Tensor Product. The Fubini-Haagerup tensor product, which we define below, is developed in the same way as the weak* Haagerup tensor product is in [5], and is general enough to give us a usable form of the extended Haagerup tensor product of [14].

Let us begin by recalling the theory of the weak* Haagerup tensor product, but in the context we require. Let $\mathcal{H}$ be a Hilbert space of Hilbertian dimension $\dim \mathcal{H}$ and I be an index set of cardinality $|I| = (\dim \mathcal{H})^2$. Squaring accommodates the case that $\dim \mathcal{H}$ is finite. Following arguments from [43, Theo. 3.1], which may also be found in [21], we have that if $S \in \mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$ then there exist $[v_i]$ in $M_{1,I}(\mathcal{B}(\mathcal{H}))$ and $[w_i]$ in $M_{I,1}(\mathcal{B}(\mathcal{H}))$ such that for any a in $\mathcal{B}(\mathcal{H})$

$$Sa = \sum_{i \in I} v_i a w_i = \begin{bmatrix} v_{i_1} & v_{i_2} & \cdots \end{bmatrix} a \begin{bmatrix} w_{i_1} \\ w_{i_2} \\ \vdots \end{bmatrix}$$

where we well-order $I = \{i_1, i_2, \dots\}$ to represent matrix multiplication, and, moreover

$$\|S\|_{cb} = \|[v_i]\| \|[w_i]\| = \left\| \sum_{i \in I} v_i v_i^* \right\|^{1/2} \left\| \sum_{i \in I} w_i^* w_i \right\|^{1/2}.$$

The size of the index set I may be deduced by tracing through the proofs the representation theorem for completely bounded maps found in [36, Theo. 2.7] and of Stinespring's Theorem [46]; or see [16, Sec. 5.2] for these results together. The aforementioned arguments can be adapted to see that if $[S_{ij}] \in M_n(\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))) \cong \mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}), M_n(\mathcal{B}(\mathcal{H})))$ then there are matrices $[v_{i\iota}]$ in $M_{n,I}(\mathcal{B}(\mathcal{H}))$ and $[w_{\iota j}]$ in $M_{I,n}(\mathcal{B}(\mathcal{H}))$ such that

$$S_{ij}a = \sum_{\iota \in I} v_{i\iota} a w_{\iota j} \quad \text{and} \quad \|S\|_{cb} = \|[v_{i\iota}]\| \|[w_{\iota j}]\|.$$

Then, following [5], we define

$$(2.1) \quad \mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H}) = \left\{ \sum_{i \in I} v_i \otimes w_i : \begin{array}{l} [v_i] \in M_{1,I}(\mathcal{B}(\mathcal{H})) \text{ and} \\ [w_i] \in M_{I,1}(\mathcal{B}(\mathcal{H})) \end{array} \right\}.$$

Here, two sums of elementary tensors, $\sum_{i \in I} v_i \otimes w_i$ and $\sum_{i \in I} v'_i \otimes w'_i$ will be regarded to be the same if $\sum_{i \in I} v_i a w_i = \sum_{i \in I} v'_i a w'_i$ for every a in $\mathcal{B}(\mathcal{H})$. The norm on $M_n(\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H}))$ is given by

$$(2.2) \quad \|u\|_{w^*h} = \min \left\{ \|v\| \|w\| : \begin{array}{l} v \in M_{n,I}(\mathcal{B}(\mathcal{H})), \ w \in M_{I,n}(\mathcal{B}(\mathcal{H})) \\ \text{and } u = v \odot w \end{array} \right\}$$

where for $v = [v_{i\ell}]$ in $M_{n,I}(\mathcal{B}(\mathcal{H}))$ and $w = [w_{\ell j}]$ in $M_{I,n}(\mathcal{B}(\mathcal{H}))$, $v \odot w$ has ij th component given by

$$(v \odot w)_{ij} = \sum_{\ell \in I} v_{i\ell} \otimes w_{\ell j}.$$

If $u = \sum_{i \in I} v_i \otimes w_i$ in $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$, define $T_u : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ by

$$(2.3) \quad T_u a = \sum_{i \in I} v_i a w_i.$$

Then, by our construction of $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$, $u \mapsto T_u$ is a surjective complete isometry from $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$ to $\mathcal{CB}^\sigma(\mathcal{B}(\mathcal{H}))$. As in [5], we have that $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$ is a dual operator space. We also note that it can be deduced from [43] that $\mathcal{B}(\mathcal{H}) \otimes^h \mathcal{B}(\mathcal{H})$ imbeds completely isometrically into $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$.

The *left* and *right slice maps*, L_F and R_F below, will be our major tool for identifying useful subspaces of $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$.

Proposition 2.1. *If $F \in \mathcal{B}(\mathcal{H})^*$ and $u = \sum_{i \in I} v_i \otimes w_i$ in $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$, the sums*

$$(2.4) \quad L_F u = \sum_{i \in I} F(v_i) w_i \quad \text{and} \quad R_F u = \sum_{i \in I} F(w_i) v_i.$$

*converge in norm. Moreover, L_F and R_F define completely bounded maps from $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$ to $\mathcal{B}(\mathcal{H})$.*

Proof. We will show this only for L_F .

To see that $\sum_{i \in I} F(v_i) w_i$ converges in norm, we simply note that since F is automatically completely bounded we have that $F^{(1,I)}[v_i] = [F(v_i)] \in M_{1,I}$, where $M_{1,I}$ is the row Hilbert space $\ell^2(I)_r$, so the coefficients $F(v_i)$ are square summable.

To see that L_F defines a completely bounded linear map, we must first check that it is well-defined. If ξ, η and ζ are vectors in $\mathcal{H}$, let $\omega_{\xi, \eta}$ denote the usual vector functional and $\zeta \otimes \eta^*$ the usual rank 1 operator. If

$[v_i]$ in $M_{1,I}(\mathcal{B}(\mathcal{H}))$ and $[w_i]$ in $M_{I,1}(\mathcal{B}(\mathcal{H}))$ are such that $\sum_{i \in I} v_i a w_i = 0$ for every a in $\mathcal{B}(\mathcal{H})$, then, letting $u = \sum_{i \in I} v_i \otimes w_i$ we have that

$$[R_{\omega_{\xi,\eta}}] \zeta = \sum_{i \in I} \langle w_i \xi | \eta \rangle v_i \zeta = \sum_{i \in I} \langle \xi | w_i^* \eta \rangle v_i \zeta = \sum_{i \in I} v_i (\zeta \otimes \eta^*) w_i \xi = 0 \cdot \xi = 0$$

from which it follows that $R_{\omega_{\xi,\eta}} u = 0$. Then, we have that $\omega_{\xi,\eta}(L_F u) = F(R_{\omega_{\xi,\eta}} u) = 0$ for each ξ and η in $\mathcal{H}$, so $L_F u = 0$.

The complete boundedness of L_F is now easy to check. Indeed, if $u = v \odot w$ in $M_n(\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H}))$, as in (2.2), with $\|u\|_{w^*h} = \|v\| \|w\|$, then

$$\|L_F^{(n)} u\| = \left\| \left(F^{(n,I)} v \right) w \right\| \leq \|F\| \|v\| \|w\| = \|F\| \|u\|_{w^*h}$$

so $\|L_F\|_{cb} \leq \|F\|$.

(Thanks go to D. Blecher for pointing out to the author that it is not *a priori* clear that L_F and R_F are well-defined. The proof here uses the idea of [5, Prop. 3.7 (i)].) $\square$

Now let $\mathcal{V}$ and $\mathcal{W}$ be closed subspaces of $\mathcal{B}(\mathcal{H})$. We define the *Fubini-Haagerup tensor product* by

$$\mathcal{V} \otimes^{fh} \mathcal{W} = \left\{ u \in \mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H}) : \begin{array}{l} L_F u \in \mathcal{W} \text{ and } R_F u \in \mathcal{V} \\ \text{for each } F \text{ in } \mathcal{B}(\mathcal{H})^* \end{array} \right\}.$$

It is clear that $\mathcal{V} \otimes^{fh} \mathcal{W}$ is a norm-closed subspace of $\mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$. It follows from the theorem below that $\mathcal{V} \otimes^{fh} \mathcal{W}$ can be defined for abstract operator spaces, independently of their completely isometric representations in $\mathcal{B}(\mathcal{H})$.

Theorem 2.2. *Let $\mathcal{V}$ and $\mathcal{W}$ be closed subspaces of $\mathcal{B}(\mathcal{H})$ and $u \in \mathcal{B}(\mathcal{H}) \otimes^{w^*h} \mathcal{B}(\mathcal{H})$. Then the following are equivalent:*

- (i) $u \in \mathcal{V} \otimes^{fh} \mathcal{W}$.
- (ii) $R_\omega u \in \mathcal{V}$ and $L_\omega u \in \mathcal{W}$ for each ω in $\mathcal{B}(\mathcal{H})_*$.
- (iii) There exist v in $M_{1,I}(\mathcal{V})$ and w in $M_{I,1}(\mathcal{W})$ such that

$$u = v \odot w \quad \text{and} \quad \|u\|_{w^*h} = \|v\| \|w\|.$$

In particular, we have that if $u \in M_n(\mathcal{V} \otimes^{fh} \mathcal{W})$, then

$$(2.5) \quad \|u\|_{w^*h} = \min \left\{ \|v\| \|w\| : \begin{array}{l} v \in M_{n,I}(\mathcal{V}), w \in M_{I,n}(\mathcal{W}) \\ \text{and } u = v \odot w \end{array} \right\}.$$

Proof. That (i) implies (ii) is trivial. If (ii) holds, then we have that $R_\omega u \in \mathcal{V}$ and $L_\omega u \in \mathcal{W}$ for each vector functional $\omega = \omega_{\xi,\eta}$. Then the proof follows identically that of [43, Theo. 4.1], adapted to accommodate uncountable index sets; we observe that the definition of *strong independence* [43, Def. 3.2] is suitable for uncountable column matrices of operators. Finally, if (iii) holds then (i) follows from Proposition 2.1. $\square$

Corollary 2.3. *If $\mathcal{V}, \mathcal{W}, \mathcal{X}$ and $\mathcal{Y}$ are all operator spaces, $S \in \mathcal{CB}(\mathcal{V}, \mathcal{X})$ and $T \in \mathcal{CB}(\mathcal{W}, \mathcal{Y})$, then the map $S \otimes T : \mathcal{V} \otimes \mathcal{W} \rightarrow \mathcal{X} \otimes \mathcal{Y}$ extends to a completely bounded map, again denoted $S \otimes T$, from $\mathcal{V} \otimes^{fh} \mathcal{W}$ to $\mathcal{X} \otimes^{fh} \mathcal{Y}$. If S and T are each complete isometries, then so too is $S \otimes T$.*

Proof. It should first be checked that $S \otimes T$ is well-defined on $\mathcal{V} \otimes^{fh} \mathcal{W}$. If $u = 0$ in $\mathcal{V} \otimes^{fh} \mathcal{W}$, then we must have that $(S \otimes T)u = 0$ since for each F in $\mathcal{X}^*$ we have that $[L_{F^\circ}(S \otimes T)]u = TL_{F \circ S}u = T0 = 0$, where the fact that $L_{F \circ S}u = 0$ follows from Proposition 2.1 and the Arveson-Wittstock Extension Theorem. Similarly, $[R_{F'^\circ}(S \otimes T)]u = 0$ for each F' in $\mathcal{Y}^*$. It then follows easily from (2.2) that $\|S \otimes T\|_{cb} \leq \|S\|_{cb} \|T\|_{cb}$.

That $S \otimes T$ is a complete isometry when S and T each are, also follows from (2.2). $\square$

It is also immediate from Theorem 2.2 that if $\mathcal{V}$ and $\mathcal{W}$ are dual operator spaces then $\mathcal{V} \otimes^{fh} \mathcal{W} = \mathcal{V} \otimes^{w^*h} \mathcal{W}$, the *weak* Haagerup tensor product*; see [5, Theo. 3.1]. Hereafter we prefer to use the notation $\otimes^{w^*h}$ in this case. As in [5, Theo. 3.2], if $\mathcal{V}$ and $\mathcal{W}$ have operator preduals $\mathcal{V}_*$ and $\mathcal{W}_*$, then $\mathcal{V} \otimes^{w^*h} \mathcal{W} \cong (\mathcal{V}_* \otimes^h \mathcal{W}_*)^*$ via the dual pairing

$$(2.6) \quad \left\langle \sum_{i \in I} v_i \otimes w_i, \nu \otimes \omega \right\rangle = \sum_{i \in I} \nu(v_i) \omega(w_i).$$

If $\mathcal{V}$ and $\mathcal{W}$ are any pair of operator spaces, the *extended Haagerup tensor product* is defined in [14] by

$$\mathcal{V} \otimes^{eh} \mathcal{W} = \left\{ B \in (\mathcal{V}^* \otimes^h \mathcal{W}^*)^* : \begin{array}{l} f \mapsto B(f \otimes g_0) \text{ and } g \mapsto B(f_0 \otimes g) \\ \text{are weak}^* \text{ continuous for any} \\ g_0 \text{ in } \mathcal{W}^* \text{ and } f_0 \text{ in } \mathcal{V}^* \end{array} \right\}$$

Theorem 2.4. *For any pair of operator spaces $\mathcal{V}$ and $\mathcal{W}$, $\mathcal{V} \otimes^{eh} \mathcal{W} \cong \mathcal{V} \otimes^{fh} \mathcal{W}$ completely isometrically.*

Proof. Both $\mathcal{V} \otimes^{eh} \mathcal{W}$ and $\mathcal{V} \otimes^{fh} \mathcal{W}$ naturally inject completely isometrically into $(\mathcal{V}^* \otimes^h \mathcal{W}^*)^* \cong \mathcal{V}^{**} \otimes^{w^*h} \mathcal{W}^{**}$, with the same range. $\square$

We will now dispense with the notation $\mathcal{V} \otimes^{fh} \mathcal{W}$, preferring to use $\mathcal{V} \otimes^{eh} \mathcal{W}$ in its place, unless both of $\mathcal{V}$ and $\mathcal{W}$ are dual spaces.

3. SCHUR MULTIPLIERS

3.1. Classical Schur Multipliers. We motivate our study in Subsection 3.2 by exposing the theory of classical Schur multipliers in our context.

Let us first fix an index set I and consider the Hilbert space $\ell^2(I)$. It will be convenient to identify $\mathcal{B}(\ell^2(I))$ with $M_{I,I}$, the space of $I \times I$ scalar matrices representing bounded operators on $\ell^2(I)$, with respect to the canonical ortho-normal basis $\{\delta_i\}_{i \in I}$.

A function $u : I \times I \rightarrow \mathbb{C}$ is called a *Schur multiplier* if $[u(i, j)\alpha_{ij}] \in M_{I,I}$ for every $[\alpha_{ij}]$ in $M_{I,I}$. An application of the Closed Graph Theorem shows that the operator

$$(3.1) \quad S_u : M_{I,I} \rightarrow M_{I,I} \quad \text{given by} \quad S_u[\alpha_{ij}] = [u(i, j)\alpha_{ij}]$$

is bounded. We denote the space of Schur multipliers by $V^\infty(I)$ and let $\|u\|_{V^\infty} = \|S_u\|$ for each u in $V^\infty(I)$.

The von Neumann algebra $\ell^\infty(I)$ of bounded functions on I can be naturally identified with $\{M_\varphi : \varphi \in \ell^\infty(I)\}$, the algebra of multiplication operators on $\ell^2(I)$: if $\varphi \in \ell^\infty(I)$, then $M_\varphi \delta_i = \varphi(i)\delta_i$. We thus identify $\ell^\infty(I)$ with the algebra of diagonal matrices in $M_{I,I}$.

In the notation of the previous section, we have that

$$\ell^\infty(I) \otimes^{w^*h} \ell^\infty(I) = \left\{ \sum_{k \in K} \varphi_k \otimes \psi_k : \left\| \sum_{k \in K} |\varphi_k|^2 \right\|_\infty^{1/2} \left\| \sum_{k \in K} |\psi_k|^2 \right\|_\infty^{1/2} < +\infty \right\}.$$

Here, K is an index set chosen so that $|K| = |I|^2$, where the squaring accommodates the case that I is finite. We know from Theorem 2.2 that this space has norm given by

$$\|v\|_{w^*h} = \min \left\{ \left\| \sum_{k \in K} |\varphi_k|^2 \right\|_\infty^{1/2} \left\| \sum_{k \in K} |\psi_k|^2 \right\|_\infty^{1/2} : v = \sum_{k \in K} \varphi_k \otimes \psi_k \right\}.$$

We have, then, that the operator $T_v : M_{I,I} \rightarrow M_{I,I}$ takes the form

$$(3.2) \quad T_v[\alpha_{ij}] = \sum_{k \in K} \text{diag}(\varphi_k) [\alpha_{ij}] \text{diag}(\psi_k)$$

where $\text{diag}(\varphi)$ is the diagonal matrix with diagonal entries $\text{diag}(\varphi)_{ii} = \varphi(i)$.

The following result is actually [37, Prop. 1.1], but with a different proof.

Theorem 3.1. *A function $u : I \times I \rightarrow \mathbb{C}$ is in $V^\infty(I)$ if and only if there is a $v = \sum_{k \in K} \varphi_k \otimes \psi_k$ in $\ell^\infty(I) \otimes^{w^*h} \ell^\infty(I)$ such that $u = v$, i.e.*

$$u(i, j) = \sum_{k \in K} \varphi_k(i) \psi_k(j)$$

for each (i, j) in $I \times I$. Moreover, $S_u = T_v$ in this case, and hence $\|u\|_{V^\infty} = \|v\|_{w^*h}$.

Proof. First, if $v = \sum_{k \in K} \varphi_k \otimes \psi_k$ in $\ell^\infty(I) \otimes^{w^*h} \ell^\infty(I)$, write

$$v(i, j) = \sum_{k \in K} \varphi_k(i) \psi_k(j)$$

for each (i, j) in $I \times I$. Then if $[\alpha_{ij}] \in M_{I,I}^{\text{fin}}$, i.e. $[\alpha_{ij}]$ has only finitely many non-zero entries, it follows from (3.2) that

$$(3.3) \quad T_v[\alpha_{ij}] = \sum_{k \in K} [\varphi_k(i) \alpha_{ij} \psi_k(j)] = \left[\sum_{k \in K} \varphi_k(i) \alpha_{ij} \psi_k(j) \right] = [v(i, j) \alpha_{ij}].$$

Since T_v is weak*-weak* continuous on $M_{I,I}$ and $M_{I,I}^{\text{fin}}$ is weak*-dense in $M_{I,I}$, (3.3) holds for any $[\alpha_{ij}]$ in $M_{I,I}$. In particular, we see that v , qua function on $I \times I$, is in $V^\infty(I)$ and with $T_v = S_v$.

Now if $u \in V^\infty(I)$, let $\check{u}(i, j) = u(j, i)$. Then $\|\check{u}\|_{V^\infty} = \|u\|_{V^\infty}$. Indeed the transpose map $x \mapsto x^t$ is an isometry on $M_{I,I}$ and $S_{\check{u}} = (S_u x^t)^t$ for each x . Let $T_{I,I}$ denote the $I \times I$ trace class matrices, so $T_{I,I}^* \cong M_{I,I}$ via $\langle t, x \rangle = \text{tr}(tx)$. We note that $S_{\check{u}}(T_{I,I}^{\text{fin}}) \subset T_{I,I}^{\text{fin}}$ and that $S_u = (S_{\check{u}}|_{T_{I,I}^{\text{fin}}})^*$. This implies that $S_{\check{u}}$ is bounded on $T_{I,I}^{\text{fin}}$ and hence extends to $T_{I,I}$, with $S_u = (S_{\check{u}}|_{T_{I,I}})^*$. Hence S_u is weak*-weak* continuous on $M_{I,I}$. Moreover, S_u is clearly an $\ell^\infty(I)$ -bimodule map. Since $\ell^\infty(I)$ is locally cyclic on $\ell^2(I)$ (and, in fact, cyclic if I is countable), S_u is completely bounded by [43, Sec. 2]. Hence we find that $S_u = T_v$ for some v in $\ell^\infty(I) \otimes^{w^*h} \ell^\infty(I)$, by [43, Theo. 3.1]. It then follows from above that $v = u$ and that $\|v\|_{w^*h} = \|T_v\| = \|S_u\| = \|u\|_{V^\infty}$. $\square$

We can use Theorem 3.1 to deduce the following well-known result of Grothendieck. See [37, Prop. 1.1] or [38, Theo. 5.1], for example.

Corollary 3.2. *A function $u : I \times I \rightarrow \mathbb{C}$ is in $V^\infty(I)$ if and only if there is a Hilbert space $\mathcal{H}$, and bounded functions $\xi, \eta : I \rightarrow \mathcal{H}$ such that*

$$u(i, j) = \langle \xi(i) | \eta(j) \rangle$$

for each (i, j) in $I \times I$. We write $u = u_{\xi, \eta}$, in this case. Moreover, $\|u\|_{V^\infty} = \inf\{\|\xi\|_\infty \|\eta\|_\infty : u = u_{\xi, \eta}\}$.

Proof. If $u \in V^\infty(I)$, then $u \in \ell^\infty(I) \otimes^{w^*h} \ell^\infty(I)$ and we may write $u = \sum_{k \in K} \varphi_k \otimes \psi_k$ where $\|u\|_{V^\infty} = \|\sum_{k \in K} |\varphi_k|^2\|_\infty^{1/2} \|\sum_{k \in K} |\psi_k|^2\|_\infty^{1/2}$. Let $\xi, \eta : I \rightarrow \ell^2(K)$ be given by

$$(3.4) \quad \xi(i) = (\varphi_k(i))_{k \in K} \quad \text{and} \quad \eta(i) = \left(\overline{\psi_k(i)} \right)_{k \in K}.$$

Then $\|\xi\|_\infty = \|\sum_{k \in K} |\varphi_k|^2\|_\infty^{1/2}$ and $\|\eta\|_\infty = \|\sum_{k \in K} |\psi_k|^2\|_\infty^{1/2}$, and $u = u_{\xi, \eta}$.

Conversely, let $u = u_{\xi, \eta}$ where $\xi, \eta : I \rightarrow \mathcal{H}$ are bounded functions. Let $\{e_k\}_{k \in K}$ be an ortho-normal basis for $\mathcal{H}$. Then for each k in K , let $\varphi_k, \psi_k : I \rightarrow \mathbb{C}$ be given by

$$\varphi_k(i) = \langle \xi(i) | e_k \rangle \quad \text{and} \quad \psi_k(i) = \langle e_k | \eta(i) \rangle.$$

Then it follows from Parseval's formula that $\sum_{k \in K} |\varphi_k(i)|^2 = \|\xi(i)\|^2$, so $\|\sum_{k \in K} |\varphi_k|^2\|_\infty^{1/2} = \|\xi\|_\infty$. Similarly, $\|\sum_{k \in K} |\psi_k|^2\|_\infty^{1/2} = \|\eta\|_\infty$ and we get that $u = \sum_{k \in K} \varphi_k \otimes \psi_k$. $\square$

3.2. Measurable Schur Multipliers. For this subsection we let (X, μ) denote a measure space, with positive measure μ , which satisfies the Radon-Nikodym Theorem so that $L^1(X, \mu)^* \cong L^\infty(X, \mu)$. This is satisfied whenever μ is σ -finite or if μ is a regular measure on the Borel sets of a locally compact space X (see [27, III.12] for the latter). We note that since (X, μ) is not necessarily σ -finite, elements of $L^\infty(X, \mu)$ can be identified as functions only up to locally null sets. A subset N of X is called μ -*locally null* if $\mu(N \cap E) = 0$ for any μ -measurable subset E of X such that $\mu(E) < +\infty$. Thus the norm on $L^\infty(X, \mu)$ is given by

$$\begin{aligned} \|\varphi\|_\infty &= \text{ess sup}_{x \in X} |\varphi(x)| \\ &= \inf \{c \in \mathbb{R}^+ : \{x \in X : |\varphi(x)| > c\} \text{ is locally null}\}. \end{aligned}$$

We will want spaces of Hilbert space valued L^∞ -functions. For our purposes it is sufficient to consider only weakly measurable such functions in the sense of [10]. We outline the construction of such spaces to clarify our notation. If $\mathcal{H}$ is a Hilbert space, let $L_{\mathcal{H}}^\infty(X, \mu)_0$ denote the set of functions $\xi : X \rightarrow \mathcal{H}$ such that

- (i) for all η in $\mathcal{H}$, $x \mapsto \langle \xi(x) | \eta \rangle$ is measurable, and
- (ii) $\|\xi\|_\infty = \text{ess sup}_{x \in X} \|\xi(x)\| < +\infty$.

Note that if $\mathcal{H}$ is separable, then (ii) follows from

- (ii') $\sup\{\|\langle \xi(\cdot) | \eta \rangle\|_\infty : \eta \in b_1(\mathcal{H})\} < +\infty$

by a straightforward application of the Uniform Boundedness Principle.

It is evident that $L_{\mathcal{H}}^\infty(X, \mu)_0$ is a linear space with semi-norm $\|\cdot\|_\infty$. Hence if $L_{\mathcal{H}}^\infty(X, \mu)$ denotes the quotient space $L_{\mathcal{H}}^\infty(X, \mu)_0 / \{\xi : \|\xi\|_\infty = 0\}$, it is a normed space, which can be verified to be complete. As is the case with $L^\infty(X, \mu)$, we will consider elements of $L_{\mathcal{H}}^\infty(X, \mu)$ as functions, defined μ -locally almost everywhere.

A subset E of $X \times X$ is called $\mu \times \mu$ -*locally marginally null*, if $E \subset (N_1 \times X) \cup (X \times N_2)$ for some locally null sets N_1 and N_2 in X . We will be interested in describing spaces of functions defined only up to locally marginally null sets. We note that if X is σ -finite for μ , then every $\mu \times \mu$ -locally marginally null is a $\mu \times \mu$ -null set. However, the converse does not hold: consider $X = [0, 1]$, the unit interval in $\mathbb{R}$, with μ being Lebesgue measure and $E = \{(s, s) : s \in [0, 1]\}$. For convenience, we will abbreviate “(locally) marginally almost every” to (l.)m.a.e. in the sequel.

If $\xi, \eta \in L_{\mathcal{H}}^\infty(X, \mu)$, define $u_{\xi, \eta}$ for $\mu \times \mu$ -l.m.a.e. (x, y) in $X \times X$ by

$$(3.5) \quad u_{\xi, \eta}(x, y) = \langle \xi(x) | \eta(y) \rangle.$$

Note that if $\xi = \xi'$ and $\eta = \eta'$, except on μ -locally null sets N_ξ and N_η , then $u_{\xi, \eta} = u_{\xi', \eta'}$ except on $(N_\xi \times X) \cup (X \times N_\eta)$, so there is no ambiguity in defining $u_{\xi, \eta}$. Let

$$(3.6) \quad V^\infty(X, \mu) = \{u_{\xi, \eta} : \xi, \eta \in L_{\mathcal{H}}^\infty(X, \mu) \text{ for some Hilbert space } \mathcal{H}\}.$$

By using Hilbert space direct sums and tensor products, it is easy to see that $V^\infty(X, \mu)$ is an algebra. In fact, it can be further verified that it is a

Banach algebra under the norm

$$(3.7) \quad \|u\|_{V^\infty} = \inf\{\|\xi\|_\infty \|\eta\|_\infty : u = u_{\xi,\eta}\}.$$

for u in $V^\infty(X, \mu)$. However, since this fact follows from Theorem 3.3 *infra*, we will not prove it directly.

For $p = 1, 2$ or ∞ , let $L^p = L^p(X, \mu)$ below.

The space $T(X, \mu) = L^2 \otimes_\gamma L^2$ is the predual of $\mathcal{B}(L^2)$ via the identification

$$(3.8) \quad \langle S, f \otimes g \rangle = \langle Sf | \bar{g} \rangle = \int_X Sf(x)g(x)d\mu(x).$$

Elements of $T(X, \mu)$ may be regarded as functions, defined up to marginally null sets: if $\omega = \sum_{k=1}^\infty f_k \otimes g_k \in T(X, \mu)$, then let

$$(3.9) \quad \omega(x, y) = \sum_{k=1}^\infty f_k(y)g_k(x)$$

for m.a.e. (x, y) in $X \times X$. See [2, 2.2.7]. Note that the order of x and y above is purposeful, and leads to technical simplifications in the sequel. A function $u : X \times X \rightarrow \mathbb{C}$ is said to be a *multiplier* of $T(X, \mu)$ if

$$(3.10) \quad m_u \omega(x, y) = u(x, y) f(y) g(x)$$

defines an element of $T(X, \mu)$ for each elementary tensor $\omega = f \otimes g$ in $L^2 \otimes_\gamma L^2$ and does so in such a way that $\|m_u f \otimes g\|_\gamma \leq C \|f\|_2 \|g\|_2$ for a fixed constant $C > 0$. Hence m_u extends to a bounded linear map on $T(X, \mu)$. It is easy to see that two multipliers u and u' satisfy $m_u = m_{u'}$ if and only if $u = u'$ locally marginally almost everywhere. The linear space of $(\mu \times \mu$ -l.m.a.e. equivalence classes of) such functions will be denoted $MT(X, \mu)$.

Let

$$(3.11) \quad \Gamma_2(L^1, L^\infty) = \left\{ T : L^1 \rightarrow L^\infty : \begin{array}{l} T = T_2 \circ T_1 \text{ where } T_1 \in \mathcal{B}(L^1, \mathcal{H}) \\ \text{and } T_2 \in \mathcal{B}(\mathcal{H}, L^\infty), \\ \text{for some Hilbert space } \mathcal{H} \end{array} \right\}.$$

The norm on $\Gamma_2(L^1, L^\infty)$ is given by

$$\|T\|_{\Gamma_2} = \inf\{\|T_2\| \|T_1\| : T = T_2 \circ T_1 \text{ as in (3.11)}\}.$$

We note that since $L^1 = \max L^1$ and $L^\infty = \min L^\infty$, $\Gamma_2(L^1, L^\infty) = \Gamma_2^c(L^1, L^\infty)$ with the same norm, where $\Gamma_2^c(L^1, L^\infty)$ is the space of operators which factor as completely bounded maps through a column space. See [16, 13.3] or [15]. Hence, by [15, Theo. 5.3] $\Gamma_2(L^1, L^\infty) \cong (L^1 \otimes^h L^1)^*$ via

$$(3.12) \quad \langle T, f \otimes g \rangle = \langle Tf, g \rangle = \int_X Tf(x)g(x)d\mu(x)$$

and thus $\Gamma_2(L^1, L^\infty)$ attains the dual operator space structure. We note that the matrix norms on $\Gamma_2(L^1, L^\infty)$ may be described independently of this duality (see [16, 13.3]).

Since (X, μ) is assumed to satisfy the Radon-Nikodym Theorem, we have that $(L^1)^* \cong L^\infty$. Moreover, we have a natural identification of L^∞ with $\{M_\varphi : \varphi \in L^\infty\}$, the maximal Abelian subalgebra of $\mathcal{B}(L^2)$ consisting of multiplication operators, where $M_\varphi f = \varphi f$ for f in L^2 . Thus, following (2.1), we obtain the space

$$L^\infty \otimes^{w^*h} L^\infty = \left\{ \sum_{i \in I} \varphi_i \otimes \psi_i : \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_\infty \left\| \sum_{i \in I} |\psi_i|^2 \right\|_\infty < +\infty \right\}$$

with norm

$$\|u\|_{V^\infty} = \min \left\{ \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_\infty^{1/2} \left\| \sum_{i \in I} |\psi_i|^2 \right\|_\infty^{1/2} : u = \sum_{i \in I} \varphi_i \otimes \psi_i \right\}.$$

It will be convenient to represent the duality $(L^1 \otimes^h L^1)^* \cong L^\infty \otimes^{w^*h} L^\infty$ given in (2.6):

$$(3.13) \quad \left\langle \sum_{i \in I} \varphi_i \otimes \psi_i, f \otimes g \right\rangle = \sum_{i \in I} \left(\int_X \varphi_i f d\mu \right) \left(\int_X \psi_i g d\mu \right)$$

for f, g in L^1 . By [5, Cor. 3.8], $L^\infty \otimes^{w^*h} L^\infty$ injects naturally into $L^\infty \overline{\otimes} L^\infty \cong L^\infty(X \times X, \mu \times \mu)$, so we may regard $L^\infty \otimes^{w^*h} L^\infty$ as a space of measurable functions on $X \times X$. As we shall see in Theorem 3.3 *infra*, $u = u'$ in $L^\infty \otimes^{w^*h} L^\infty$ if and only if $u = u'$ locally marginally almost everywhere.

We let

$$(3.14) \quad \mathcal{B}_{L^\infty}^\sigma(\mathcal{B}(L^2)) = \left\{ T \in \mathcal{B}^\sigma(\mathcal{B}(L^2)) : \begin{array}{l} T(M_\varphi a M_\psi) = M_\varphi T(a) M_\psi \\ \text{for each } a \text{ in } \mathcal{B}(L^2) \\ \text{and } \varphi, \psi \text{ in } L^\infty \end{array} \right\}.$$

We can now state the main result of this section. This establishes a framework which will be useful in the sequel.

Theorem 3.3. *If (X, μ) is a measure space satisfying the Radon-Nikodym Theorem, then the following spaces coincide isometrically:*

- (i) $V^\infty(X, \mu)$
- (ii) $L^\infty(X, \mu) \otimes^{w^*h} L^\infty(X, \mu)$
- (iii) $MT(X, \mu)$.

Moreover, the space in (ii) is naturally completely isometrically isomorphic to each of the following spaces:

- (iv) $\mathcal{B}_{L^\infty(X, \mu)}^\sigma(\mathcal{B}(L^2(X, \mu)))$
- (v) $\Gamma_2(L^1(X, \mu), L^\infty(X, \mu))$.

*Finally, if (X, μ) is σ -finite, then any element in $L^\infty(X, \mu) \otimes^{w^*h} L^\infty(X, \mu)$ can be represented as a countable sum of elementary tensors, and the definition of $V^\infty(X, \mu)$ can be made using separable Hilbert spaces only.*

We note that if μ is counting measure on X , then $V^\infty(X, \mu) = V^\infty(X)$, so this result generalises Theorem 3.1 and Corollary 3.2. Henceforth we will

identify $V^\infty(X, \mu)$ with $L^\infty(X, \mu) \otimes^{w^*h} L^\infty(X, \mu)$, and call elements of this space *measurable Schur multipliers*.

In the case that (X, μ) is σ -finite, U. Haagerup appears to obtain the equivalence of (i), (ii) and (iii), also with countable index sets in the definition on $L^\infty(X, \mu) \otimes^{w^*h} L^\infty(X, \mu)$, and separable Hilbert spaces in the definition of $V^\infty(X, \mu)$. See [21] or [52].

Proof. As above we let L^p denote $L^p(X, \mu)$ for $p = 1, 2$ or ∞ .

(i)=(ii) This follows the proof of Corollary 3.2 almost exactly. However, if $u = \sum_{i \in I} \varphi_i \otimes \psi_i$ in $L^\infty \otimes^{w^*h} L^\infty$, we want to ensure that the analogues of (3.4) are weakly measurable. Indeed, let $\xi, \eta : X \rightarrow \ell^2(I)$ be given by

$$(3.15) \quad \xi(x) = (\varphi_i(x))_{i \in I} \quad \text{and} \quad \eta(x) = \left(\overline{\psi_i(x)} \right)_{i \in I}$$

for locally almost every x in X . Then if $\zeta = (\zeta_i)_{i \in I}$ in $\ell^2(I)$, we have that $\zeta_i = 0$ for all but countably many i and hence $\langle \xi(\cdot) | \zeta \rangle = \sum_{i \in I} \overline{\zeta_i} \varphi_i$ is measurable. Similarly η is weakly measurable.

(ii) $\cong$ (iv) The map (2.3) restricts to a surjective complete isometry from $L^\infty \otimes^{w^*h} L^\infty$ to $\mathcal{CB}_{L^\infty}^\sigma(\mathcal{B}(L^2))$ by Corollary 2.3 and [43, Theo. 3.1]. Then $\mathcal{CB}_{L^\infty}^\sigma(\mathcal{B}(L^2)) = \mathcal{B}_{L^\infty}^\sigma(\mathcal{B}(L^2))$ by [43, Theo. 3.1], since L^∞ is locally cyclic on L^2 .

(ii) $\cong$ (v) From (3.13), $L^\infty \otimes^{w^*h} L^\infty \cong (L^1 \otimes^h L^1)^*$, completely isometrically. Also, $(L^1 \otimes^h L^1)^* \cong \Gamma_2(L^1, L^\infty)$ via (3.12). We note that the complete isomorphism $L^\infty \otimes^{w^*h} L^\infty \cong \Gamma_2(L^1, L^\infty)$ is given by $u \mapsto T(u)$, where for f in L^1 , $T(u)f = L_f u$, where L_f is the left slice map given in (2.4). Indeed, if $u = \sum_{i \in I} \varphi_i \otimes \psi_i$, the factorization $T = T_2 \circ T_1$, as in (3.11), with $\mathcal{H} = \ell^2(I)$, is given by

$$T_1 f = (\langle f, \varphi_i \rangle)_{i \in I} \quad \text{and} \quad T_2 \zeta = \sum_{i \in I} \zeta_i \psi_i$$

where $\zeta = (\zeta_i)_{i \in I}$.

(ii)=(iii) (via (ii) $\cong$ (iv)) Let $u \in L^\infty \otimes^{w^*h} L^\infty$, with $u = \sum_{i \in I} \varphi_i \otimes \psi_i$ and $\|u\|_{w^*h} = \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_\infty^{1/2} \left\| \sum_{i \in I} |\psi_i|^2 \right\|_\infty^{1/2}$. It suffices to show that $m_u f \otimes g \in T(X, \mu)$ with $\|m_u f \otimes g\|_\gamma \leq \|u\|_{w^*h} \|f\|_2 \|g\|_2$ for any f, g in L^2 , to establish that $u \in \text{MT}(X, \mu)$ with $\|m_u\| \leq \|u\|_{w^*h}$. Using the definition of the projective tensor product and then Hölder's Inequality we have that

$$\begin{aligned} \|m_u f \otimes g\|_\gamma &= \left\| \sum_{i \in I} M_{\psi_i} f \otimes M_{\varphi_i} g \right\|_\gamma \leq \sum_{i \in I} \|M_{\psi_i} f\|_2 \|M_{\varphi_i} g\|_2 \\ &\leq \left(\sum_{i \in I} \|M_{\psi_i} f\|_2^2 \right)^{1/2} \left(\sum_{i \in I} \|M_{\varphi_i} g\|_2^2 \right)^{1/2}. \end{aligned}$$

Then, using Tonelli's Theorem, we have that

$$\begin{aligned}
 \sum_{i \in I} \|M_{\psi_i} f\|_2^2 &= \sum_{i \in I} \int_X |\psi_i f|^2 d\mu = \int_X \sum_{i \in I} |\psi_i f|^2 d\mu \\
 (3.16) \quad &= \int_X \sum_{i \in I} |\psi_i|^2 |f|^2 d\mu \leq \left\| \sum_{i \in I} |\psi_i|^2 \right\|_\infty \|f\|_2^2
 \end{aligned}$$

and, similarly, $\sum_{i \in I} \|M_{\varphi_i} g\|_2^2 \leq \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_\infty \|g\|_2^2$. Hence

$$\|m_u f \otimes g\|_\gamma \leq \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_\infty^{1/2} \left\| \sum_{i \in I} |\psi_i|^2 \right\|_\infty^{1/2} \|f\|_2 \|g\|_2 = \|u\|_{w^*h} \|f\|_2 \|g\|_2$$

as required.

Now suppose that $u \in \text{MT}(X, \mu)$. For φ and ψ in L^∞ , the adjoint of $M_\psi \otimes M_\varphi$, qua operator on $T(X, \mu)$, in the dual pairing described in (3.8), is $T_{\varphi \otimes \psi}$. Indeed we have for a in $\mathcal{B}(L^2)$ and f, g in L^2 ,

$$\begin{aligned}
 \langle a, M_\psi \otimes M_\varphi(f \otimes g) \rangle &= \langle a, M_\psi f \otimes M_\varphi g \rangle = \langle a M_\psi f | M_\varphi^* \bar{g} \rangle \\
 &= \langle M_\varphi a M_\psi f | g \rangle = \langle T_{\varphi \otimes \psi} a, f \otimes g \rangle.
 \end{aligned}$$

Since $M_\psi \otimes M_\varphi m_u = m_u M_\psi \otimes M_\varphi$ for all φ and ψ in L^∞ , $T_{\varphi \otimes \psi} m_u^* = m_u^* T_{\varphi \otimes \psi}$. Hence $m_u^* \in \mathcal{B}_{L^\infty}^\sigma(\mathcal{B}(L^2))$ and there is u' in $L^\infty \otimes^{w^*h} L^\infty$ such that $m_u^* = T_{u'}$. But then $m_{u'} = m_u$ so $u' = u$ locally marginally almost everywhere, and hence $u \in L^\infty \otimes^{w^*h} L^\infty$.

We finally deal with the case where (X, μ) is σ -finite. We can decompose $X = \bigcup_{n \in \mathbb{N}} E_n$ where each E_n is measurable with $\mu(E_n) < +\infty$. Then, given $u = \sum_{i \in I} \varphi_i \otimes \psi_i$ in $L^\infty \otimes^{w^*h} L^\infty$, using (3.16) we have

$$\sum_{i \in I} \int_X |\psi_i \chi_{E_n}|^2 d\mu \leq \left\| \sum_{i \in I} |\psi_i|^2 \right\|_\infty \mu(E_n)$$

so $\psi_i \chi_{E_n} = 0$ (almost everywhere) for all but countably many indices i . It then follows that $\psi_i \neq 0$ for at most countably many indices i . The same holds for the functions φ_i . Hence if $u \in L^\infty \otimes^{w^*h} L^\infty$, we can write $u = \sum_{i=1}^\infty \varphi_i \otimes \psi_i$. Following (3.15), it is clear that only separable Hilbert spaces are needed in the definition of $V^\infty(X, \mu)$.

(The author would like to express his gratitude to L. Turowska for suggesting the proof of (ii)=(iv), given here. This is a generalised version of the proof of [45, Prop. 4.1].) $\square$

We observe that, in general, we can bound the size of I by $|I| \leq \aleph_0 \cdot c(X, \mu)$, where $\aleph_0$ is the first infinite cardinal and $c(X, \mu)$ is the smallest cardinality of a cover of X by μ -finite sets.

We close this section with an asymmetric version of Theorem 3.3. If (X, μ) and (Y, ν) are measure spaces, each satisfying the Radon-Nikodym Theorem, then we can consider the following spaces:

$$(\mathbf{i}') \quad V^\infty(X, Y; \mu, \nu)$$

- (ii') $L^\infty(X, \mu) \otimes^{w^*h} L^\infty(Y, \nu)$
- (iii') $L^\infty(X, \mu) \mathcal{B}_{L^\infty(Y, \nu)}(\mathcal{B}(L^2(Y, \nu), L^2(X, \mu)))$ (left $L^\infty(X, \mu)$ and right $L^\infty(Y, \nu)$ -module maps)
- (iv') $\Gamma_2(L^1(X, \mu), L^\infty(Y, \nu),)$
- (v') $MT(Y, X; \nu, \mu)$, where $T(Y, X; \nu, \mu) = L^2(Y, \nu) \otimes^\gamma L^2(X, \mu)$.

The space in (i') is defined analogously to that in (3.6); note that $V^\infty(X, \mu) = V^\infty(X, X; \mu, \mu)$. Then the spaces (i')-(v') are isometrically isomorphic. Moreover, the spaces (ii'), (iii') and (iv') have natural operator space structures in which they are completely isometrically isomorphic.

That these spaces are isomorphic can be proved just as Theorem 3.3, *mutatis mutandis*. Also note that if we take the disjoint union measure space $(X \dot{\cup} Y, \mu \dot{\cup} \nu)$, then all of the spaces (i')-(v') can be taken to be “corners” of the spaces (i)-(v) of Theorem 3.3, but with the latter mentioned spaces taken over the measure space $(X \dot{\cup} Y, \mu \dot{\cup} \nu)$.

3.3. Continuous Schur Multipliers. For this subsection we will let X and Y denote locally compact Hausdorff spaces. Since it will add no extra difficulties in notation, we will state asymmetrical versions of all our results.

Let

$$(3.17) \quad V_0(X, Y) = \mathcal{C}_0(X) \otimes^h \mathcal{C}_0(Y) \quad \text{and} \quad V_b(X, Y) = \mathcal{C}_b(X) \otimes^h \mathcal{C}_b(Y).$$

These are clearly closed subalgebras of $\ell^\infty(X) \otimes^{w^*h} \ell^\infty(Y)$ and hence Banach algebras under pointwise operations. If X and Y are compact we will write $V(X, Y) = V_0(X, Y) = V_b(X, Y)$. This algebra is called the *Varopoulos algebra* in [45].

Proposition 3.4. *The Banach algebra $V_0(X, Y)$ is regular with Gel'fand spectrum $X \times Y$. The same is true for $V_b(X, Y)$, but with spectrum $bX \times bY$, where bX and bY denote the Stone-Ćech compactifications of X and Y .*

Proof. By Grothendieck's Inequality (see [9, 14.5], for example), $V_0(X, Y) = \mathcal{C}_0(X) \otimes^\gamma \mathcal{C}_0(Y)$. Since $\mathcal{C}_0(X)$ and $\mathcal{C}_0(Y)$ are regular and satisfy the metric approximation property, the proposition follows from [48]. The result for $V_b(X, Y)$ follows immediately from the identities $\mathcal{C}_b(X) \cong \mathcal{C}(bX)$ and $\mathcal{C}_b(Y) \cong \mathcal{C}(bY)$ and the reasoning above. $\square$

In the sequel we will need spaces which extend $V_0(X, Y)$ and $V_b(X, Y)$. Let

$$(3.18) \quad V^0(X, Y) = \mathcal{C}_0(X) \otimes^{eh} \mathcal{C}_0(Y) \quad \text{and} \quad V^b(X, Y) = \mathcal{C}_b(X) \otimes^{eh} \mathcal{C}_b(Y).$$

It follows from Corollary 2.3 that these spaces are closed subalgebras of $\ell^\infty(X) \otimes^{w^*h} \ell^\infty(Y)$ and hence Banach algebras under pointwise operations. Moreover, we obtain from Theorem 2.2, the following.

Proposition 3.5. *A function $u : X \times Y \rightarrow \mathbb{C}$ defines an element of $V^0(X, Y)$ (respectively $V^b(X, Y)$), if and only if there exist families of functions $\{\varphi_i\}_{i \in I}$ in $\mathcal{C}_0(X)$ (respectively $\mathcal{C}_b(X)$) and $\{\psi_i\}_{i \in I}$ in $\mathcal{C}_0(Y)$ (respectively $\mathcal{C}_b(Y)$)*

such that $u(x, y) = \sum_{i \in I} \varphi_i(x) \psi_i(y)$ for each (x, y) in $X \times Y$ (i.e. $u = \sum_{i \in I} \varphi_i \otimes \psi_i$), and

$$\left\| \sum_{i \in I} |\varphi_i|^2 \right\|_{\infty} \left\| \sum_{i \in I} |\psi_i|^2 \right\|_{\infty} < +\infty.$$

Moreover the norm of u in $V^0(X, Y)$ (respectively $V^b(X, Y)$) is then

$$\|u\|_V = \min \left\{ \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_{\infty}^{1/2} \left\| \sum_{i \in I} |\psi_i|^2 \right\|_{\infty}^{1/2} : u = \sum_{i \in I} \varphi_i \otimes \psi_i \right\}.$$

Observe that we can obtain a bound for the cardinality of the index set I : $|I| \leq \min\{d(\mathcal{C}_b(X)), d(\mathcal{C}_b(Y))\}$, where for any Banach space $\mathcal{X}$, $d(\mathcal{X})$ denotes the *density degree* of $\mathcal{X}$. If X and Y admit a σ -finite regular Borel measures μ and ν such that $\mu(U) > 0$ and $\nu(V) > 0$ for each open subset U of X and V of Y , then $V^b(X, Y)$ (and hence $V^0(X, Y)$) is a closed subspace of $V^\infty(X, Y; \mu, \nu)$, and hence I can be taken to be countable by Theorem 3.3.

We will find the following description of $V^b(X, Y)$ useful. Let for a Hilbert space $\mathcal{H}$,

$$\mathcal{C}_b(X, w^*\text{-}\mathcal{H}) = \left\{ \xi : X \rightarrow \mathcal{H} : \begin{array}{l} \xi \text{ is continuous when } \mathcal{H} \text{ has the weak}^* \\ \text{topology, and } \|\xi\|_{\infty} = \sup_{x \in X} \|\xi(x)\| < +\infty \end{array} \right\}.$$

Similarly, define $\mathcal{C}_b(Y, w^*\text{-}\mathcal{H})$. If $\xi \in \mathcal{C}_b(X, w^*\text{-}\mathcal{H})$ and $\eta \in \mathcal{C}_b(Y, w^*\text{-}\mathcal{H})$, let $u_{\xi, \eta} : X \times Y \rightarrow \mathbb{C}$ be given by $u_{\xi, \eta}(x, y) = \langle \xi(x) | \eta(y) \rangle$.

Proposition 3.6. *If X and Y are locally compact Hausdorff spaces, then $u \in V^b(X, Y)$ if and only if there are a Hilbert space $\mathcal{H}$ and functions ξ in $\mathcal{C}_b(X, w^*\text{-}\mathcal{H})$ and η in $\mathcal{C}_b(Y, w^*\text{-}\mathcal{H})$ such that $u = u_{\xi, \eta}$. Moreover, $\|u\|_V = \min \{ \|\xi\|_{\infty} \|\eta\|_{\infty} : u = u_{\xi, \eta} \text{ as above} \}$.*

Proof. This follows the proof of Corollary 3.2 almost exactly. However, if $u = \sum_{i \in I} \varphi_i \otimes \psi_i$ in $V^b(X, Y)$, we want to ensure that the analogues of (3.4) are suitably continuous. Indeed, let $\xi : X \rightarrow \ell^2(I)$ and $\eta : Y \rightarrow \ell^2(I)$ be given by

$$\xi(x) = (\varphi_i(x))_{i \in I} \quad \text{and} \quad \eta(y) = \left(\overline{\psi_i(y)} \right)_{i \in I}$$

for each x in X and y in Y . To see that $\xi \in \mathcal{C}_b(X, w^*\text{-}\mathcal{H})$, let $\zeta = (\zeta_i)_{i \in I}$ in $\ell^2(I)$. Then we have that

$$|\langle \xi(\cdot) | \zeta \rangle| = \left| \sum_{i \in I} \zeta_i \varphi_i \right| \leq \|\zeta\|_2 \left\| \sum_{i \in I} |\varphi_i|^2 \right\|_{\infty}^{1/2}$$

so the middle series converges uniformly, and hence $\langle \xi(\cdot) | \zeta \rangle$ is continuous. Similarly, $\eta \in \mathcal{C}_b(Y, w^*\text{-}\mathcal{H})$. $\square$

Although an element u in $V^b(X, Y)$ always unambiguously describes a function on $X \times Y$, it may not describe a continuous function. This fact is pointed out in [42]. As an example, let us take $X = Y = [0, 1]$, the unit interval in $\mathbb{R}$. For each n in $\mathbb{N}$ let $\varphi_n = \psi_n : [0, 1] \rightarrow [0, 1]$ be a continuous function supported on $[\frac{1}{2^{n+1}}, \frac{1}{2^n}]$ such that $\varphi_n(\frac{3}{2^{n+2}}) = \psi_n(\frac{3}{2^{n+2}}) = 1$ (where $\frac{3}{2^{n+2}}$ is simply the midpoint of $\frac{1}{2^{n+1}}$ and $\frac{1}{2^n}$). Then it is clear that

$$\left\| \sum_{n=1}^{\infty} |\varphi_n|^2 \right\|_{\infty} \left\| \sum_{n=1}^{\infty} |\psi_n|^2 \right\|_{\infty} = 1 \cdot 1 = 1$$

and hence that $u = \sum_{n=1}^{\infty} \varphi_n \otimes \psi_n \in V^b([0, 1], [0, 1])$ with norm 1. However, while $u(0, 0) = 0$, $u(\frac{3}{2^{n+2}}, \frac{3}{2^{n+2}}) = 1$ for each n , and hence u is not continuous at 0. We note, then, that for any choice of spaces X and Y , finding the Gel'fand spectrum for $V^b(X, Y)$ or $V^0(X, Y)$, should prove to be challenging.

It would be interesting to have intrinsic descriptions of $V^b(X, Y) \cap \mathcal{C}_b(X \times Y)$, $V^0(X, Y) \cap \mathcal{C}_b(X \times Y)$ and $V^0(X, Y) \cap \mathcal{C}_0(X \times Y)$.

4. COMPLETELY BOUNDED MULTIPLIERS OF THE FOURIER ALGEBRA

For this section the symbol $\mathcal{H}$, possibly with subscript, will denote a Hilbert space, and $\mathcal{U}(\mathcal{H})$ the group of unitaries on $\mathcal{H}$, topologized with the weak operator topology.

Our standard references for harmonic analysis are [27], [17] and [1].

Let G be a locally compact group with left Haar integral $f \mapsto \int_G f(s) ds$. For $p = 1, 2$, let $L^p(G)$ denote the space of (almost everywhere equivalence classes of) functions f for which $\|f\|_p = (\int_G |f(s)|^p ds)^{1/p} < +\infty$. Then $L^1(G)$ is an involutive Banach algebra under the product of convolution, $f * g(t) = \int_G f(s) g(s^{-1}t) ds$ for almost every t , and with involution $f^*(t) = \frac{1}{\Delta(t)} \overline{f(t^{-1})}$, where the overline denotes complex conjugation and $\Delta : G \rightarrow \mathbb{R}^{>0}$ is the Haar modular function.

If π is a continuous unitary representation of G , i.e. a continuous homomorphism from G into $\mathcal{U}(\mathcal{H}_\pi)$, then it forms a canonical non-degenerate involutive representation $\pi_1 : L^1(G) \rightarrow \mathcal{B}(\mathcal{H}_\pi)$ via the weak*-converging integrals $\pi_1(f) = \int_G f(s) \pi(s) ds$. We let C_π^* be the norm closure of $\pi_1(L^1(G))$ in $\mathcal{B}(\mathcal{H}_\pi)$, and $C^*(G)$ denote the *enveloping C^* -algebra* of G , i.e. the enveloping C^* -algebra of $L^1(G)$.

The *Fourier* and *Fourier-Stieltjes algebras*, $A(G)$ and $B(G)$, are defined in [17]. We recall that $B(G)$ is the space of matrix coefficients of all continuous unitary representations of G ; i.e. the space of functions of the form $s \mapsto \langle \pi(s) \xi | \eta \rangle$ where π is a continuous unitary representation of G on $\mathcal{H}_\pi$, and $\xi, \eta \in \mathcal{H}_\pi$. $B(G)$ is the dual of $C^*(G)$ via $\langle a, \langle \pi(\cdot) \xi | \eta \rangle \rangle = \langle \pi_*(a) \xi | \eta \rangle$, where $\pi_* : C^*(G) \rightarrow \mathcal{B}(\mathcal{H}_\pi)$ is the representation induced by π , i.e. by π_1 . We will refer to $\sigma(B(G), C^*(G))$ as *the weak* topology* on $B(G)$, although its uniqueness is not assured. It is well-known that $B(G)$ is a commutative

Banach algebra (under pointwise operations) and $A(G)$ is the closed ideal in $B(G)$ generated by compactly supported matrix coefficients.

If π is a continuous unitary representation of G , let A_π be the norm closure of $\text{span}\{\langle \pi(\cdot)\xi|\eta\rangle : \xi, \eta \in \mathcal{H}_\pi\}$ in $B(G)$. Then, by [1, 2.2], $A_\pi^* \cong \text{VN}_\pi$, where VN_π is the von Neumann algebra generated by $\pi(G)$. Thus if ϖ denotes the universal representation of G , we have $B(G) = A_\varpi$. The algebra $W^*(G) = \text{VN}_\varpi$ is called the *enveloping von Neumann algebra* of G . Also, $A(G) = A_\lambda$ where λ is the left regular representation of G . We let $\text{VN}(G) = \text{VN}_\lambda$, and call it the *group von Neumann algebra*. We will also let B_π denote the weak*-closure of A_π in $B(G)$. By standard functional analysis, $B_\pi \cong (C_\pi^*)^*$. In particular, we let $B_r(G) = B_\lambda$. Then $B_r(G)$ is an ideal in $B(G)$, and is the dual of the *reduced group C*-algebra* $C_r^*(G) = C_\lambda^*$.

4.1. Multipliers and Completely Bounded Multipliers. A *completely contractive Banach algebra* is a Banach algebra $\mathcal{A}$, equipped with an operator space structure such that the multiplication map $m : \mathcal{A} \otimes_\wedge \mathcal{A} \rightarrow \mathcal{A}$ is a complete contraction, and hence extends to a complete contraction $m : \hat{\mathcal{A}} \otimes \mathcal{A} \rightarrow \mathcal{A}$.

The Fourier-Stieltjes algebra $B(G)$, as the dual of $C^*(G)$, is a completely contractive Banach algebra. Indeed, the multiplication map $m : B(G) \otimes_\wedge B(G) \rightarrow B(G)$ has adjoint $m^* : W^*(G) \rightarrow W^*(G) \overline{\otimes} W^*(G)$, given for s in G by $m^*\varpi(s) = \varpi(s) \otimes \varpi(s)$. By the universal property of $W^*(G)$, m^* is a *-homomorphism and hence a complete contraction; thus so too is m .

Any of the subspaces A_π of $B(G)$ obtains the same operator space structure qua subspace of $B(G)$ as it does as the predual of VN_π . Hence if A_π is a subalgebra of $B(G)$, then it is a completely contractive Banach algebra. In particular, $A(G)$ is a completely contractive Banach algebra.

Let $\mathcal{A}$ be a semi-simple commutative Banach algebra with spectrum X , or more generally, a subalgebra of $\mathcal{C}_b(X)$ for which there is some a in $\mathcal{A}$ such that $a(x) \neq 0$ for each x in X , and comes equipped with a norm under which it is a Banach algebra. The *multipliers* of $\mathcal{A}$ are given by

$$M\mathcal{A} = \{u : X \rightarrow \mathbb{C} : ua \in \mathcal{A} \text{ for each } a \text{ in } \mathcal{A}\}.$$

An application of the Closed Graph Theorem shows that the map $m_u : \mathcal{A} \rightarrow \mathcal{A}$, given by

$$(4.1) \quad m_u a = ua$$

is bounded. Moreover, it can be shown that $\{m_u : u \in M\mathcal{A}\}$ is the strong operator topology closed subalgebra of $\mathcal{A}$ -module maps in $\mathcal{B}(\mathcal{A})$. See [32], for example. If $\mathcal{A}$ has an operator space structure let

$$M_{cb}\mathcal{A} = \{u \in M\mathcal{A} : m_u \in \mathcal{CB}(\mathcal{A})\}.$$

The operator space structure is given as follows: for $[u_{ij}]$ in $M_n(M_{cb}\mathcal{A})$ let

$$\|[u_{ij}]\|_{cbm} = \left\| m_{[u_{ij}]} \right\|_{cb}$$

where $m_{[u_{ij}]}$ in $\mathcal{CB}(\mathcal{A}, M_n(\mathcal{A}))$ is given by

$$(4.2) \quad m_{[u_{ij}]}a = [m_{u_{ij}}a] = [u_{ij}a]$$

for a in $\mathcal{A}$.

If $\mathcal{A}$ is, in addition to the properties assumed above, a completely contractive Banach algebra, then it injects completely contractively into $M_{cb}\mathcal{A}$. Moreover, if $\mathcal{A}$ has a contractive bounded approximate identity $\{e_\alpha\}$, then the injection $\mathcal{A} \hookrightarrow M_{cb}\mathcal{A}$ is a complete isometry. Indeed, if $[a_{ij}] \in M_n(\mathcal{A})$ then

$$(4.3) \quad \|[a_{ij}]\| \geq \|[a_{ij}]\|_{cbm} \geq \lim_\alpha \|[a_{ij}e_\alpha]\| = \|[a_{ij}]\|.$$

Now let G be a locally compact group and $A(G)$ the Fourier algebra of G . The following Proposition can be proved exactly as [8, Prop. 1.2], but using the fact that the adjoint map $T \mapsto T^*$ from $\mathcal{CB}(\mathcal{V})$ to $\mathcal{CB}^\sigma(\mathcal{V}^*)$ is a complete isometry for any operator space $\mathcal{V}$, and that the restriction map $T \mapsto T|_{\mathcal{V}_0}$ from $\mathcal{CB}(\mathcal{V})$ to $\mathcal{CB}(\mathcal{V}_0, \mathcal{V})$, for any closed subspace $\mathcal{V}_0$, is a complete contraction

Proposition 4.1. *If $u_{ij} : G \rightarrow \mathbb{C}$ for $i, j = 1, \dots, n$, then the following are equivalent:*

- (i) $[u_{ij}] \in b_1(M_n(M_{cb}A(G)))$.
 - (ii) *The operator defined by $\lambda(s) \mapsto [u_{ij}(s)\lambda(s)]$, for s in G , extends to a weakly continuous complete contraction, $M_{[u_{ij}]} : VN(G) \rightarrow M_n(VN(G))$.*
 - (iii) *The operator defined by $\lambda_1(f) \mapsto [\lambda_1(u_{ij}f)]$, for f in $L^1(G)$, extends to a complete contraction $\tilde{M}_{[u_{ij}]} : C_r^*(G) \rightarrow M_n(C_r^*(G))$.*
 - (iv) $[u_{ij}] \in b_1(M_n(M_{cb}B_r(G)))$.
- In (ii), $M_{[u_{ij}]}T = [M_{u_{ij}}T]$, for T in $VN(G)$, where for $i, j = 1, \dots, n$, $M_{u_{ij}} \in \mathcal{CB}^\sigma(VN(G))$ is the adjoint of $m_{u_{ij}}$ in $\mathcal{CB}(A(G))$.*

Following (iii) of the above proposition, we may consider $M_{cb}A(G)$ to be the “multipliers of $C_r^*(G)$ ”. We will see below that $B(G)$ represents the multipliers of the enveloping C^* -algebra, $C^*(G)$. In [51] and [52], it is shown that $B(G)$ imbeds as a maximal Abelian subalgebra of $\mathcal{CB}(C^*(G))$. We prove this below, and also show that the operator space structure, induced on $B(G)$ as a subspace of $\mathcal{CB}(C^*(G))$, is the same as the operator space structure induced on $B(G)$ as the dual of $C^*(G)$.

Proposition 4.2. *If $u_{ij} : G \rightarrow \mathbb{C}$ for $i, j = 1, \dots, n$, then the following are equivalent:*

- (i) $[u_{ij}] \in b_1(M_n(B(G)))$.
- (ii) $[u_{ij}] \in b_1(M_n(M_{cb}B(G)))$.
- (iii) *The operator defined by $\varpi(s) \mapsto [u_{ij}(s)\varpi(s)]$, for s in G , extends to a weakly continuous complete contraction $M_{[u_{ij}]} : W^*(G) \rightarrow M_n(W^*(G))$.*
- (iv) *The operator defined by $\varpi_1(f) \mapsto [\varpi_1(u_{ij}f)]$, for f in $L^1(G)$, extends to a complete contraction $\tilde{M}_{[u_{ij}]} : C^*(G) \rightarrow M_n(C^*(G))$.*

Thus $B(G)$ is completely isometrically isomorphic to the maximal Abelian subalgebra $\{\tilde{M}_u : u \in B(G)\}$ of $\mathcal{CB}(C^*(G))$.

Proof. The fact that (i) is equivalent to (ii) follows (4.3). That (ii), (iii) and (iv) are equivalent can be shown exactly as the equivalence of (ii)-(iv) in the preceding proposition, i.e. by following [8, Prop. 1.2].

We have that $\{\tilde{M}_u : u \in B(G)\}$ is a maximal Abelian subalgebra of $\mathcal{CB}(C^*(G))$, since if T in $\mathcal{CB}(C^*(G))$ commutes with every $\tilde{M}_u$, where $u \in B(G)$, then T^* is a $B(G)$ -module map on $B(G)$, and hence $T = m_u$ for some u in $B(G)$. $\square$

We note that, in fact, $\{\tilde{M}_u : u \in B(G)\}$ is a maximal Abelian subalgebra of $\mathcal{B}(C^*(G))$.

We can refine the well known result that $B(G) \subset M_{cb}A(G)$ in general, and that $B(G) = M_{cb}A(G)$ when G is amenable. See [8, Cor. 1.8] and [33], for example.

Corollary 4.3. $B(G) \subset M_{cb}A(G)$, and the identity injection $B(G) \hookrightarrow M_{cb}A(G)$ is a complete contraction. If G is amenable, then ι is a completely isometric isomorphism.

Proof. If $u \in B(G)$ let m_u be the multiplier operator in $\mathcal{CB}(B(G))$. Then, $m_u A(G) \subset A(G)$, since $A(G)$ is an ideal in $B(G)$. Moreover, $m_u \mapsto m_u|_{A(G)}$ is a complete contraction.

If G is amenable, then $B(G) = B_r(G)$, completely isometrically. the result then follows from Proposition 4.1 (iv) and Proposition 4.2 (ii). $\square$

5. INVARIANT SCHUR MULTIPLIERS

For this section, let G be a locally compact group with left Haar measure m . Let $V^\infty(G) = V^\infty(G, m)$. Hence, if G is discrete, $V^\infty(G)$ is just the space of usual Schur multipliers on $\mathcal{B}(\ell^2(G))$.

Let

$$(5.1) \quad V_{\text{inv}}^\infty(G) = \left\{ u \in V^\infty(G) : \begin{array}{l} \text{for all } r \text{ in } G, u(sr, t) = u(s, tr^{-1}) \\ \text{for l.m.a.e. } (s, t) \text{ in } G \times G \end{array} \right\}$$

In the identification $V^\infty(G) = L^\infty(G) \otimes^{w^*h} L^\infty(G) \cong (L^1(G) \otimes^h L^1(G))^*$ in Theorem 3.3 we have that

$$V_{\text{inv}}^\infty(G) = \bigcap_{r \in G} \ker (\text{id}_{L^\infty(G)} \otimes \rho(r^{-1})^* - \rho(r)^* \otimes \text{id}_{L^\infty(G)}).$$

Here $\rho : G \rightarrow \mathcal{B}(L^1(G))$ is the right regular representation on $L^1(G)$, given by $\rho(r)f(s) = \Delta(r)f(sr)$, for r in G , f in $L^1(G)$ and almost all s in G . Hence $V_{\text{inv}}^\infty(G)$ is a weak* closed subspace of $V^\infty(G)$.

If G is a discrete group, then $V^\infty(G)$ is a space of functions, rather than a space of equivalence classes of functions. Thus for u in $V_{\text{inv}}^\infty(G)$ it is evident that we can define a function $u_G : G \rightarrow \mathbb{C}$ such that

$$u_G(st^{-1}) = u(s, t)$$

for all (s, t) in $G \times G$. For a non-discrete group this is more subtle.

Lemma 5.1. *Suppose G is a non-discrete locally compact group, and let $u \in V_{\text{inv}}^\infty(G)$. Consider u as a function on $G \times G$, i.e. as a representative of its equivalence class. Then there is a locally null subset E_u of G such that for t in $G \setminus E_u$, $u(s, t^{-1}s) = u(e, t^{-1})$ for locally almost every s in G .*

Proof. Let for t in G , $F_t = \{s \in G : u(s, t^{-1}s) \neq u(e, t^{-1})\}$. Then let $E_u = \{t \in G : F_t \text{ is not locally null}\}$. We will show that E_u is locally null. If $t \in E_u$ and $s \in F_t$, then $u(s, t^{-1}s) \neq u(e, t^{-1}) = u(ss^{-1}, t^{-1})$, so there is an r in G , namely $r = s^{-1}$, such that $u(sr, t^{-1}) \neq u(s, t^{-1}r^{-1})$. Hence

$$\bigcup_{t \in E_u} F_t \times \{t^{-1}\} \subset \{(s, t) \in G \times G : u(sr, t^{-1}) \neq u(s, t^{-1}r^{-1}) \text{ for some } r \text{ in } G\}$$

and thus $\bigcup_{t \in E_u} F_t \times \{t^{-1}\}$ is locally marginally null. Let N_1 and N_2 be locally null subsets of G such that

$$\bigcup_{t \in E_u} F_t \times \{t^{-1}\} \subset (N_1 \times G) \cup (G \times N_2).$$

Note that $F_t \setminus N_1 \neq \emptyset$ for each t in E_u , since, for such t , F_t is not locally null. Then, since $(\bigcup_{t \in E_u} (F_t \setminus N_1) \times \{t^{-1}\}) \cap (N_1 \times G) = \emptyset$, we have that

$$\bigcup_{t \in E_u} (F_t \setminus N_1) \times \{t^{-1}\} \subset G \times N_2.$$

Hence $E_u^{-1} \subset N_2$ so $E_u \subset N_2^{-1}$, and thus E_u is locally null. $\square$

If $u \in V_{\text{inv}}^\infty(G)$, considered as a function as in the above lemma, but bounded, define for t in G ,

$$(5.2) \quad u_G(t) = \begin{cases} u(e, t^{-1}) & \text{if } t \in G \setminus E_u \\ \lim_{s \in V, V \in \mathcal{V}_t} u(e, s^{-1}) & \text{if } t \in E_u \end{cases}$$

where for t in E_u , $\mathcal{V}_t$ is any fixed ultra-filter over $G \setminus E_u$ containing the sets $U \setminus E_u$, where U is a neighbourhood of t . Note that $U \setminus E_u$ is always non-empty, since an m -locally null set in G never contains an open set. It is shown in the next theorem that u_G is dependent only on the choice of u in $V_{\text{inv}}^\infty(G)$ qua equivalence class. Moreover, it is shown that u_G is automatically continuous.

Let $T(G) = L^2(G) \otimes^\gamma L^2(G)$, as in Subsection 3.2. Also recall the convention that for $\omega = f \otimes g$ in $T(G)$, $\omega(s, t) = f(t)g(s)$ for m.a.e. (s, t) in $G \times G$.

Theorem 5.2. *For each u in $V_{\text{inv}}^\infty(G)$, $u_G \in M_{cb}A(G)$. The map $u \mapsto u_G$ induces a complete contraction from $V_{\text{inv}}^\infty(G)$ to $M_{cb}A(G)$.*

Proof. The map $P : T(G) \rightarrow A(G)$ given by

$$Pf \otimes g(t) = \langle \lambda(t)f | \overline{g} \rangle = \int_G f(t^{-1}s)g(s)ds$$

for t in G and f, g in $L^2(G)$, is a surjective contraction whose adjoint, P^* , is the usual imbedding $VN(G) \hookrightarrow \mathcal{B}(L^2(G))$. Thus if $u \in V_{\text{inv}}^\infty(G)$, letting m_u be as in (3.10), we have that $P(m_u\omega) \in A(G)$ for ω in $T(G)$. If G is not discrete, let E_u be as the above lemma; if G is discrete let $E_u = \emptyset$. Then for t in $G \setminus E_u$ we have

$$P(m_u\omega)(t) = \int_G u(s, t^{-1}s)\omega(s, t^{-1}s)ds = u_G(t)P\omega(t).$$

Let $v \in A(G)$ with $v = P\omega$, for some ω in $T(G)$. Since v is continuous, using the notation of (5.2), we have that

$$u_G(t)v(t) = \lim_{s \in V, V \in \mathcal{V}_t} u_G(s)v(s) = \lim_{s \in V, V \in \mathcal{V}_t} P(m_u\omega)(s) = P(m_u\omega)(t)$$

for t in E_u . Hence $u_G(t)v(t) = P(m_u\omega)(t)$ for all t in G , so $u_G v \in A(G)$. Then, by definition, $u_G \in MA(G)$ with $P \circ m_u = m_{u_G} \circ P$, where m_{u_G} in $\mathcal{B}(A(G))$ is as defined in (4.1). Taking adjoints we have that $T_u \circ P^* = (P \circ m_u)^* = (m_{u_G} \circ P)^* = P^* \circ M_{u_G}$, where M_{u_G} in $\mathcal{B}^\sigma(VN(G))$ is the adjoint of m_{u_G} , and hence for x in $VN(G)$ we have that,

$$T_u x = m_u^* x = M_{u_G} x.$$

Thus $M_{u_G} = T_u|_{VN(G)}$, so $\|u_G\|_{cbm} = \|M_{u_G}\|_{cb} \leq \|T_u\|_{cb} = \|u\|_{V^\infty}$. It also follows that for $[u_{ij}]$ in $M_n(V_{\text{inv}}^\infty(G))$, $\|[u_{ij}]\|_{cbm} \leq \|[u_{ij}]_G\|_{V^\infty}$. $\square$

The following theorem refines the main result in [7]. We consider this to be the centrepiece of this paper. Our proof uses modern operator space theoretic techniques, and gives us a complete isometry. Also, it exposes the relationship between $M_{cb}A(G)$ and $L^\infty(G)$ -module maps on $\mathcal{B}(L^2(G))$ via Theorem 3.3. Many of our methods are motivated by the elegant exposition found in [22].

Theorem 5.3. *If $u_{ij} : G \rightarrow \mathbb{C}$ are continuous functions for $i, j = 1, \dots, n$, then the following are equivalent:*

- (i) $[u_{ij}] \in b_1(M_n(M_{cb}A(G)))$.
- (ii) *For any finite subset F of G , if $u_{ij}^F(s, t) = u_{ij}(st^{-1})$ for s, t in F , then $[u_{ij}^F] \in b_1(M_n(V^\infty(F)))$.*
- (iii) *If u_{ij}^G is given by $u_{ij}^G(s, t) = u_{ij}(st^{-1})$ for s, t in G , then $[u_{ij}^G] \in b_1(M_n(V_{\text{inv}}^\infty(G_d)))$.*
- (iv) $[u_{ij}^G] \in b_1(M_n(V_{\text{inv}}^\infty(G)))$.

In particular, the map $u \mapsto u^G$ is a complete isometry from $M_{cb}A(G)$ onto $V_{\text{inv}}^\infty(G)$.

Proof. (i) $\Rightarrow$ (ii) Let $F = \{s_1, \dots, s_m\}$, and for $k, l = 1, \dots, m$ let E_{kl} in $M_m(VN(G))$ be identified by

$$E_{kl} \cong \lambda(s_k s_l^{-1}) \otimes e_{kl} \quad \text{in} \quad VN(G) \bar{\otimes} M_m \cong M_m(VN(G))$$

where $\{e_{kl}\}_{k,l=1,\dots,m}$ is the standard matrix unit for M_m . Then $\{E_{kl}\}_{k,l=1,\dots,m}$ is itself a matrix unit so $\mathcal{N}_F = \text{span}\{E_{kl}\}_{k,l=1,\dots,m} \cong M_m$ via $E_{kl} \mapsto e_{kl}$. Let us identify $M_m \cong \mathcal{B}(\ell^2(F))$. Note that for v in $V^\infty(F)$, $S_v e_{kl} = v(s_k, s_l) e_{kl}$, for each k, l , where S_v is defined as in (3.1). If $[v_{ij}] \in M_n(V^\infty(F))$, let $S_{[v_{ij}]}x = [S_{v_{ij}}x]$ for x in $\mathcal{B}(\ell^2(F))$. We then have

$$\begin{aligned} M_{[u_{ij}]}^{(m)} E_{kl} &\cong [M_{u_{ij}} \lambda(s_k s_l^{-1}) \otimes e_{kl}] = [u_{ij}(s_k s_l^{-1}) \lambda(s_k s_l^{-1}) \otimes e_{kl}] \\ &\cong [u_{ij}(s_k s_l^{-1}) E_{kl}] \cong [u_{ij}^F(s_k, s_l) e_{kl}] \cong S_{[u_{ij}^F]} e_{kl} \end{aligned}$$

so $[u_{ij}^F] \in M_n(V^\infty(F))$. Moreover,

$$\|[u_{ij}^F]\|_{V^\infty} = \|S_{[u_{ij}^F]}\|_{cb} = \|M_{[u_{ij}]}|_{\mathcal{N}_F}\|_{cb} \leq \|[u_{ij}]\|_{cbm} \leq 1$$

as required.

(ii) $\Rightarrow$ (iii) Let $\mathcal{F}$ denote the collection of all finite subsets of G , directed by inclusion. We have for each F in $\mathcal{F}$, a natural completely isometric imbedding

$$V^\infty(F) = \ell^\infty(F) \otimes^h \ell^\infty(F) \subset \ell^\infty(G_d) \otimes^{w*h} \ell^\infty(G_d) = V^\infty(G_d)$$

and hence an isometric imbedding $M_n(V^\infty(F)) \subset M_n(V^\infty(G_d))$. Denote the image of $[u_{ij}^F]$ under this imbedding again by $[u_{ij}^F]$. Since $M_n(V^\infty(G_d))$ is a dual space (with predual $T_n(\ell^1(G_d) \otimes^h \ell^1(G_d))$), the net $\left([u_{ij}^F]\right)_{F \in \mathcal{F}}$ must have a weak* cluster point in $b_1(M_n(V^\infty(G_d)))$. Using the fact that $\bigcup_{F \in \mathcal{F}} \ell^1(F) \otimes^h \ell^1(F)$ is dense in $\ell^1(G_d) \otimes^h \ell^1(G_d)$, it is easy to see that

$$[u_{ij}^G] = w^* - \lim_{F \in \mathcal{F}} [u_{ij}^F]$$

so $[u_{ij}^G] \in b_1(M_n(V^\infty(G_d)))$. That $[u_{ij}^G] \in M_n(V_{\text{inv}}^\infty(G_d))$ is clear.

(iii) $\Rightarrow$ (iv) By (2.2) we may write $[u_{ij}^G]$ as a matricial product $A \odot B$, where $A \in M_{n,I}(\ell^\infty(G_d))$ and $B \in M_{I,n}(\ell^\infty(G_d))$, for some index set I , with $\|A\| \|B\| \leq 1$. We may explicitly write this down: well-order $I = \{\iota_1, \iota_2, \dots\}$ so that we have

$$(5.3) \quad [u_{ij}^G] = \begin{bmatrix} \varphi_{1\iota_1} & \varphi_{1\iota_2} & \varphi_{1\iota_3} & \cdots \\ \vdots & \vdots & \vdots & \\ \varphi_{n\iota_1} & \varphi_{n\iota_2} & \varphi_{n\iota_3} & \cdots \end{bmatrix} \odot \begin{bmatrix} \psi_{\iota_1 1} & \cdots & \psi_{\iota_1 n} \\ \psi_{\iota_2 1} & \cdots & \psi_{\iota_2 n} \\ \psi_{\iota_3 1} & \cdots & \psi_{\iota_3 n} \\ \vdots & & \vdots \end{bmatrix}$$

and we see that

$$u_{ij}^G = \sum_{\iota \in I} \varphi_{i\iota} \otimes \psi_{\iota j}.$$

For $i, j = 1, \dots, n$ let $\xi_i, \eta_j : G \rightarrow \ell^2(I)$ be given by

$$\xi_i(s) = (\varphi_{i\iota}(s))_{\iota \in I} \quad \text{and} \quad \eta_j(s) = \left(\overline{\psi_{\iota j}(s)}\right)_{\iota \in I}$$

so that $u_{ij}^G(s, t) = \langle \xi_i(s) | \eta_j(t) \rangle$ for s, t in G . If we consider each ξ_i and η_j as row vectors with $\ell^\infty(G_d)$ entries, i.e. as elements of $M_{1,I}(\ell^\infty(G_d))$, then we may rewrite (5.3) as

$$(5.4) \quad [u_{ij}^G] = \begin{bmatrix} \xi_1 \\ \vdots \\ \xi_n \end{bmatrix} \odot [\eta_1^* \quad \cdots \quad \eta_n^*].$$

Let $\mathcal{H}' = \overline{\text{span}}\{\eta_j(t) : j = 1, \dots, n, t \in G\}$ and p' in $\mathcal{B}(\ell^2(I))$ denote the orthogonal projection onto $\mathcal{H}'$. Then let $\mathcal{H} = \overline{\text{span}}\{p'\xi_i(s) : i = 1, \dots, n, s \in G\}$ and p in $\mathcal{B}(\ell^2(I))$ denote the orthogonal projection onto $\mathcal{H}$. Clearly $pp' = p$, and $\langle p\xi_i(s) | \eta_j(t) \rangle = \langle \xi_i(s) | \eta_j(t) \rangle$ for all indices i and j , and all s, t in G . Let $\tilde{\xi}_i, \tilde{\eta}_j : G \rightarrow \ell^2(I)$ be given by

$$\tilde{\xi}_i(s) = p\xi_i(s) \quad \text{and} \quad \tilde{\eta}_j(s) = p\eta_j(s).$$

For any $i = 1, \dots, n$ the function $\tilde{\xi}_i$ is weakly continuous. Indeed for any s in G and η in $\ell^2(I)$, $\langle \tilde{\xi}_i(s) | \eta \rangle = \langle \xi_i(s) | p\eta \rangle$ where $p\eta \in \mathcal{H}'$. Then, for any t in G and $j = 1, \dots, n$, $\langle \xi_i(s) | \eta_j(t) \rangle = u_{ij}^G(s, t) = u_{ij}(st^{-1})$, where $s \mapsto u_{ij}(st^{-1})$ is continuous by assumption. Similarly, each $\tilde{\eta}_j$ is continuous. Thus for any $i, j = 1, \dots, n$ and ι in I , the functions

$$(5.5) \quad \tilde{\varphi}_{i\iota} = \langle \tilde{\xi}_i(\cdot) | e_\iota \rangle \quad \text{and} \quad \tilde{\psi}_{\iota j} = \langle e_\iota | \tilde{\eta}_j(\cdot) \rangle$$

are continuous. Then, following the notation of (5.3), if

$$\tilde{A} = \begin{bmatrix} \tilde{\varphi}_{1\iota_1} & \tilde{\varphi}_{1\iota_2} & \tilde{\varphi}_{1\iota_3} & \cdots \\ \vdots & \vdots & \vdots & \\ \tilde{\varphi}_{n\iota_1} & \tilde{\varphi}_{n\iota_2} & \tilde{\varphi}_{n\iota_3} & \cdots \end{bmatrix} \quad \text{and} \quad \tilde{B} = \begin{bmatrix} \tilde{\psi}_{\iota_1 1} & \cdots & \tilde{\psi}_{\iota_1 n} \\ \tilde{\psi}_{\iota_2 1} & \cdots & \tilde{\psi}_{\iota_2 n} \\ \tilde{\psi}_{\iota_3 1} & \cdots & \tilde{\psi}_{\iota_3 n} \\ \vdots & & \vdots \end{bmatrix}$$

we have that

$$[u_{ij}^G] = \tilde{A} \odot \tilde{B}.$$

Indeed, for s, t in G and all indices i and j ,

$$(\tilde{A} \odot \tilde{B})_{ij}(s, t) = \langle \tilde{\xi}_i(s) | \tilde{\eta}_j(t) \rangle = \langle p\xi_i(s) | \eta_j(t) \rangle = \langle \xi_i(s) | \eta_j(t) \rangle = u_{ij}^G(s, t).$$

Let us identify $\mathcal{B}(\ell^2(I)) \cong M_{I,I}$. Then let p^t and $\bar{p}$ denote the transpose and conjugate, respectively, of the matrix p , so $\|p^t\| = \|\bar{p}\| = \|p\| \leq 1$. Identifying $M_{n,I}(\ell^\infty(G_d)) \cong \ell^\infty(G_d, M_{n,I})$, in the notation of (5.4) we have that for s in G ,

$$\tilde{A}(s) = \begin{bmatrix} \tilde{\xi}_1(s) \\ \vdots \\ \tilde{\xi}_n(s) \end{bmatrix} = \begin{bmatrix} \xi_1(s)p^t \\ \vdots \\ \xi_n(s)p^t \end{bmatrix} = \begin{bmatrix} \xi_1(s) \\ \vdots \\ \xi_n(s) \end{bmatrix} p^t = A(s)p^t.$$

Similarly, identifying $M_{n,I}(\ell^\infty(G_d)) \cong \ell^\infty(G_d, M_{n,I})$, we have that $\tilde{B}(s) = \bar{p}B(s)$. Hence $\|\tilde{A}\| \leq \|A\|$ and $\|\tilde{B}\| \leq \|B\|$, so $\|\tilde{A}\| \|\tilde{B}\| \leq 1$. Thus we see that $\left[u_{ij}^G \right] \in b_1(M_n(V_{\text{inv}}^\infty(G)))$.

(iv) $\Rightarrow$ (i) This follows from Theorem 5.2. $\square$

It is immediately evident from Theorem 5.3 that $M_{cb}A(G) = M_{cb}A(G_d) \cap \mathcal{C}_b(G)$ where G_d denotes the group G with its discrete topology. We observe this fact in a broader context in Corollary 6.3 below.

Let us consider, for the moment, the case when G is amenable, and has a left invariant mean m . We will find it notationally convenient to regard this mean as a finitely additive measure on G .

Corollary 5.4. [45, Prop. 5.2] *If G is amenable, then $V_{\text{inv}}^\infty(G) \cong B(G)$, completely isometrically, via $u \mapsto u_G$. Moreover $V_{\text{inv}}^\infty(G)$ is a completely complemented subspace of $V^\infty(G)$.*

Proof. That $M_{cb}A(G) = B(G)$, in this case, is given in Corollary 4.3. It remains to see that $V_{\text{inv}}^\infty(G)$ is a completely complemented subspace of $V^\infty(G)$. For u in $V^\infty(G)$ and r in G , let $r \cdot u(s, t) = u(sr, tr^{-1})$ for l.m.a.e. (s, t) in $G \times G$. Note that $V_{\text{inv}}^\infty(G) = \{u \in V^\infty(G) : r \cdot u = u \text{ for all } r \in G\}$. Then define $E : V^\infty(G) \rightarrow V_{\text{inv}}^\infty(G)$ by

$$Eu = w^* - \int_G r \cdot u dm(r)$$

i.e. $\langle Eu, \mu \rangle = m(r \mapsto \langle r \cdot u, \mu \rangle)$ for all μ in $L^1(G) \otimes^h L^1(G)$. Then it is easy to see that E is completely contractive. Also that $EV^\infty(G) \subset V_{\text{inv}}^\infty(G)$, and that $Eu = u$ for u in $V_{\text{inv}}^\infty(G)$, follow from the left invariance of m . $\square$

It would be interesting to determine if the (complete) complementation property above characterises amenability of G .

We now return to the case of a general locally compact group G . We will give a refinement of Theorem 5.3. Let $V^b(G) = V^b(G, G)$, where $V^b(G, G)$ is defined as in (3.18). Let

$$(5.6) \quad V_{\text{inv}}^b(G) = \{u \in V^b(G) : u(sr, t) = u(s, tr^{-1}) \text{ for } s, t, r \text{ in } G\}.$$

The following is a weak version of [7] or of [28].

Corollary 5.5. *We have that $V_{\text{inv}}^\infty(G) = V_{\text{inv}}^b(G)$. Hence $V_{\text{inv}}^b(G) = M_{cb}A(G)$ completely isometrically. Thus if $u \in M_{cb}A(G)$, then there is a Hilbert space $\mathcal{H}$ and ξ, η in $\mathcal{C}_b(G, w^* - \mathcal{H})$ such that*

$$(5.7) \quad u(st^{-1}) = \langle \xi(s) | \eta(t) \rangle$$

for all s, t in G . Moreover, $\|u\|_V = \|\xi\|_\infty \|\eta\|_\infty$.

Proof. This is immediate from the proof of (iii) $\Rightarrow$ (iv), in Theorem 5.3 above (see (5.5)), and from Proposition 3.6. $\square$

Let us finish this section by giving a second refinement of Theorem 5.3. Let $\mathcal{WAP}(G)$ denote the C^* -algebra of continuous *weakly almost periodic* functions on G . By Grothendieck's Criterion, $\mathcal{WAP}(G)$ consists of the functions φ in $\mathcal{C}_b(G)$ such that for any pair of sequences $\{s_n\}$ and $\{t_m\}$ from G , if both of the limits $\lim_n \lim_m \varphi(s_n t_m)$ and $\lim_m \lim_n \varphi(s_n t_m)$ exist, then they coincide. We note that $\mathcal{WAP}(G)$ is a closed subspace of $\mathcal{C}_b(G)$.

It is observed in [53], that $M_{cb}A(G) \subset \mathcal{WAP}(G)$. The proof is a nice application of Grothendieck's Criterion, which we used to define $\mathcal{WAP}(G)$ above, and Corollary 5.5.

Now we let

$$WV^b(G) = \mathcal{WAP}(G) \otimes^{eh} \mathcal{WAP}(G) \quad \text{and} \quad WV_{\text{inv}}^b(G) = WV^b(G) \cap V_{\text{inv}}^b(G).$$

It is clear that $WV^b(G)$ is a closed subspace of $V^b(G)$, and hence $WV_{\text{inv}}^b(G)$ is a closed subspace of $V_{\text{inv}}^b(G)$.

Corollary 5.6. *We have that $V_{\text{inv}}^\infty(G) = WV_{\text{inv}}^b(G)$. Hence $WV_{\text{inv}}^b(G) \cong M_{cb}A(G)$ completely isometrically.*

Proof. The functions $\tilde{\varphi}_{il}$ and $\tilde{\psi}_{ij}$ from (5.5) can be approximated uniformly by sums of translates of functions u_{ij} , each of which is in $M_{cb}A(G)$. Hence each $\tilde{\varphi}_{il}$ and $\tilde{\psi}_{ij}$ is weakly almost periodic. $\square$

Let us finally note that if G is compact, then

$$V_{\text{inv}}^\infty(G) = V_{\text{inv}}^b(G) \subset V(G).$$

where $V(G) = V_b(G, G) = V_0(G, G)$, as defined in (3.17). This was shown in [45].

6. APPLICATIONS

6.1. Functorial Properties of $M_{cb}A(G)$. As a first application of the results of the previous section, we will examine some of the functorial properties of the completely bounded multipliers on locally compact groups.

If X and Y are locally compact Hausdorff spaces, and $\sigma : X \rightarrow Y$ is a continuous map, let $j_\sigma : \mathcal{C}_b(Y) \rightarrow \mathcal{C}_b(X)$ be given by

$$(6.1) \quad j_\sigma \varphi = \varphi \circ \sigma.$$

It is obvious that j_σ is a $*$ -homomorphism. Moreover, if $\sigma(X)$ is dense in Y then j_σ is injective. Indeed, if $\varphi \in \ker j_\sigma$, then $\varphi(\sigma(x)) = 0$ for each x in X so $\varphi(y) = 0$ for each y in Y .

Let $V^b(X) = V^b(X, X)$ and $V^b(Y) = V^b(Y, Y)$, as in (3.18). Following Corollary 2.3 we get a completely contractive algebra homomorphism $J_\sigma = j_\sigma \otimes j_\sigma : V^b(Y) \rightarrow V^b(X)$. Moreover, J_σ is a complete isometry if $\sigma(X)$ is dense in Y .

Recall that for a locally compact group G , $V_{\text{inv}}^b(G)$ is defined in (5.6).

Proposition 6.1. *If G and H are locally compact groups and $\sigma : G \rightarrow H$ is a continuous homomorphism, then $J_\sigma V_{\text{inv}}^b(H) \subset V_{\text{inv}}^b(G)$ and J_σ is a complete isometry if $\sigma(G)$ is dense in H .*

Proof. If $u \in V_{\text{inv}}^b(H)$ then for s, r, t in G

$$J_\sigma u(sr, t) = u(\sigma(s)\sigma(r), \sigma(t)) = u(\sigma(s), \sigma(t)\sigma(r^{-1})) = J_\sigma u(s, tr^{-1})$$

so $J_\sigma u \in V_{\text{inv}}^b(G)$. That J_σ is a complete isometry if $\sigma(G)$ is dense in H follows from comments above. $\square$

We obtain the main functorial property of completely bounded multipliers by combining the above proposition with Corollary 5.5.

Theorem 6.2. *If G and H are locally compact groups and $\sigma : G \rightarrow H$ is a continuous homomorphism, then $j_\sigma M_{\text{cb}}A(H) \subset M_{\text{cb}}A(G)$, and $j_\sigma|_{M_{\text{cb}}A(H)} : M_{\text{cb}}A(H) \rightarrow M_{\text{cb}}A(G)$ defines a complete contraction, which we will again denote by j_σ . Moreover, if $\sigma(G)$ is dense in H , then j_σ is a complete isometry.*

This functorial property leads to many interesting conclusions.

Corollary 6.3. *Let G be a locally compact group.*

(i) *The inclusion $M_{\text{cb}}A(G) \hookrightarrow M_{\text{cb}}A(G_d)$ is a complete isometry, where G_d denotes the group G with discrete topology.*

(ii) *If G^{ap} denotes the almost periodic compactification of G and $\sigma_{\text{ap}} : G \rightarrow G^{\text{ap}}$ the canonical map (see [12, Sec. 6.1]), then $j_{\sigma_{\text{ap}}} : A(G^{\text{ap}}) \rightarrow M_{\text{cb}}A(G)$ is a complete isometry.*

(iii) *If H is a closed subgroup of G , then the restriction map $u \mapsto u|_H$ is a complete contraction from $M_{\text{cb}}A(G)$ to $M_{\text{cb}}A(H)$. Moreover, if H is open, then this map is a surjective complete quotient map. If H is amenable, then this map is a complete quotient map onto its range.*

(iv) *If N is a closed normal subgroup of G , and $q : G \rightarrow G/N$ is the quotient map, then $j_q : M_{\text{cb}}A(G/N) \rightarrow M_{\text{cb}}A(G)$ is a complete isometry.*

Proof. For (i), the inclusion is just the map j_{ι_d} , where $\iota_d : G_d \rightarrow G$ is the identity map. For (ii), we just note that G^{ap} is compact, so $A(G^{\text{ap}}) = B(G^{\text{ap}}) = M_{\text{cb}}A(G^{\text{ap}})$ completely isometrically, by Corollary 4.3.

For (iii), the restriction map is just j_ι , where $\iota : H \rightarrow G$ is the inclusion map. If H is open, the result follows from the equivalence of (i) and (ii) in Theorem 5.3. If H is amenable, then $j_\iota|_{B(G)} : B(G) \rightarrow B(H)$ is a complete quotient map onto its range. Indeed, the range of $j_\iota|_{B(G)}$ is $A_{\varpi|_H}$ where ϖ is the universal representation of G . The adjoint of $j_\iota|_{B(G)}$ is then the inclusion $VN_{\varpi|_H} \hookrightarrow W^*(G)$, and hence is a complete isometry. We thus obtain the factorization $j_\iota|_{B(G)} = j_\iota \circ i$, where $i : B(G) \rightarrow M_{\text{cb}}A(G)$ is the completely contractive inclusion map. Hence j_ι must be a complete quotient onto its range.

For (iv), j_q is complete isometry, since q is a surjection. $\square$

We remark that in (iv), above, the range of j_q consists of elements of $M_{cb}A(G)$ which are constant on cosets of N . In [23, p. 682] (and following that, [53, Sec. 2.2]), it is claimed that every element of $M_{cb}A(G)$ which is constant on cosets of N is in the range of j_q . If G is amenable, this result follows from that $M_{cb}A(G) = B(G)$ and [17, 2.26].

Regarding (iii), we note that it is known for an amenable group G that it is possible to have a closed subgroup H for which the restriction map from $B(G)$ to $B(H)$ is not surjective (see [17, p. 204], for example), though it is always a complete quotient onto its range. We are unable to determine if, in general, the restriction map from $M_{cb}A(G)$ to $M_{cb}A(H)$ is a (complete) quotient map onto its range.

It follows from (ii) above that if π is a finite dimensional unitary representation of G , then the imbedding $A_\pi \hookrightarrow M_{cb}A(G)$ is a complete isometry. This result is generalised below.

Corollary 6.4. *Let G be a locally compact group and π be a continuous unitary representation such that there is an amenable group H , a continuous homomorphism $\sigma : G \rightarrow H$ and a continuous unitary representation π' of H , such that $\pi = \pi' \circ \sigma$. Then the imbedding $A_\pi \hookrightarrow M_{cb}A(G)$ is a complete isometry.*

Proof. First note that $j_\sigma : B(H) \rightarrow B(G)$ is a complete contraction, since its adjoint $j_\sigma^* : W^*(G) \rightarrow W^*(H)$ is a $*$ -homomorphism, by the universal property of $W^*(G)$. Note too that each u in A_π is of the form $j_\sigma u'$ for some u' in $A_{\pi'} \subset B(H)$ by [1, Prop. 2.10], and recall that $B(H) = M_{cb}A(H)$ completely isometrically, since H is amenable. Without loss of generality we may assume that $\sigma(G)$ is dense in H , so $j_\sigma : M_{cb}A(H) \rightarrow M_{cb}A(G)$ is a complete isometry. Then if $[u_{ij}] \in M_n(A_\pi)$, we have that $[u_{ij}] = j_\sigma^{(n)} [u'_{ij}]$ for some $[u'_{ij}]$ in $M_n(B(H))$, and hence

$$\|[u_{ij}]\|_{cbm} \leq \|[u_{ij}]\| \leq \|[u'_{ij}]\| = \|[u'_{ij}]\|_{cbm} = \|j_\sigma^{(n)} [u'_{ij}]\|_{cbm} = \|[u_{ij}]\|_{cbm}.$$

Thus $\|[u_{ij}]\|_{cbm} = \|[u_{ij}]\|$. $\square$

6.2. A Predual for $M_{cb}A(G)$. In this section, we give a concrete construction of a predual of $M_{cb}A(G)$. The existence of this predual was recognised in [8], and in a more general form in [30], where it was identified as an operator predual.

Note that $\sigma : V^\infty(G) \rightarrow V^\infty(G)$ given by $\sigma u(s, t) = u(s, t^{-1})$ is a bijective complete isometry. Thus we may now consider

$$(6.2) \quad \sigma V_{\text{inv}}^\infty(G) = \left\{ u \in V^\infty(G) : \begin{array}{l} u(sr, t) = u(s, rt) \text{ for all } r \text{ in } G \\ \text{and l.m.a.e. } (s, t) \in G \times G \end{array} \right\}.$$

As in (5.2), given u in $\sigma V_{\text{inv}}^\infty(G)$, we can define a function $u_G : G \rightarrow \mathbb{C}$ such that for l.a.e. t in G

$$u_G(t) = u(s, s^{-1}t)$$

for l.a.e. s in G . Recalling (3.13), we have that $V^\infty(G) \cong (L^1(G) \otimes^h L^1(G))^*$, where the dual pairing is given by

$$\langle u, f \otimes g \rangle = \int_G \int_G u(s, t) f(s) g(t) ds dt$$

for each elementary tensor $f \otimes g$ in $L^1(G) \otimes^h L^1(G)$. If $\mathcal{X} \subset L^1(G) \otimes^h L^1(G)$, let $\mathcal{X}^\perp = \{u \in V^\infty(G) : \langle u, x \rangle = 0 \text{ for all } x \text{ in } \mathcal{X}\}$.

Let $m_0 : L^1(G) \otimes_h L^1(G) \rightarrow L^1(G)$ be the multiplication map. This is an unbounded map on an incomplete space. Let $\mathcal{K}$ be the closure of $\ker m_0$ in $L^1(G) \otimes^h L^1(G)$ and

$$(6.3) \quad Q(G) = \left(L^1(G) \otimes^h L^1(G) \right) / \mathcal{K}.$$

Endow $Q(G)$ with the quotient operator space structure.

Theorem 6.5. $\sigma V_{\text{inv}}^\infty(G) = \mathcal{K}^\perp$, and hence $Q(G)^* \cong V_{\text{inv}}^\infty(G) \cong M_{cb}A(G)$ as operator spaces.

We will call $Q(G)$ *the standard predual* of $M_{cb}A(G)$. Although its uniqueness is not assured, we will refer to the topology $\sigma(M_{cb}A(G), Q(G))$ as *the weak** topology on $M_{cb}A(G)$.

Proof. To see that $\sigma V_{\text{inv}}^\infty(G) \subset \mathcal{K}^\perp$, it suffices to see that $\sigma V_{\text{inv}}^\infty(G) \subset (\ker m_0)^\perp$. To that end, if $u \in \sigma V_{\text{inv}}^\infty(G)$ and $\mu = \sum_{k=1}^n f_k \otimes g_k$ in $\ker m_0$, then

$$\begin{aligned} \langle u, \mu \rangle &= \sum_{k=1}^n \int_G \int_G u(s, t) f_k(s) g_k(t) dt ds \\ &= \sum_{k=1}^n \int_G \int_G u(s, s^{-1}t) f_k(s) g_k(s^{-1}t) dt ds \\ &= \sum_{k=1}^n \int_G u_G(t) f_k * g_k(t) dt = \int_G u_G(t) m_0(\mu)(t) dt = 0. \end{aligned}$$

Conversely, if $u \in (\ker m_0)^\perp$, $f, g \in L^1(G)$ and δ_r is the Dirac measure at r for r in G , then $m_0(f * \delta_r \otimes g - f \otimes \delta_r * g) = 0$, so

$$\begin{aligned} 0 &= \langle u, f * \delta_r \otimes g - f \otimes \delta_r * g \rangle \\ &= \int_G \int_G u(s, t) \left(\frac{1}{\Delta(r)} f(sr^{-1})g(t) - f(s)g(r^{-1}t) \right) dt ds \\ &= \int_G \int_G (u(sr, t) - u(s, rt)) f(s)g(t) dt ds \end{aligned}$$

and hence $u(sr, t) = u(s, rt)$ for marginally almost every (s, t) in $G \times G$, so $u \in \sigma V_{\text{inv}}^\infty(G)$. Thus $(\ker m_0)^\perp \subset \sigma V_{\text{inv}}^\infty(G)$. $\square$

The map $m_0 : L^1(G) \otimes_h L^1(G) \rightarrow L^1(G)$ is surjective by Cohen's Factorization Theorem. For f in $L^1(G)$ let

$$\|f\|_Q = \inf \{ \|\mu\|_h : \mu \in L^1(G) \otimes_h L^1(G) \text{ and } m_0(\mu) = f \}.$$

The function $\|\cdot\|_Q$ is clearly a semi-norm on $L^1(G)$. Since $L^1(G)$ has a contractive bounded approximate identity, the Cohen-Hewitt Factorization Theorem [26, (32.22)] gives for any $\varepsilon > 0$ that we can write $f = g*h$, where $\|g\|_1 \leq 1$ and $\|f - h\|_1 < \varepsilon$, so that $\|g\|_1 \|h\|_1 < \|f\|_1 + \varepsilon$. Hence $\|f\|_Q \leq \|f\|_1$. Let $m : L^1(G) \otimes^h L^1(G) \rightarrow Q(G)$ be the quotient map. If $f = g*h$ in $L^1(G)$, let $i_Q(f) = m(g \otimes h)$. Then $i_Q : L^1(G) \rightarrow Q(G)$ is clearly well-defined and linear with $\|i_Q(f)\| = \|f\|_Q$.

The following shows that we may consider $Q(G)$ to be the completion of $L^1(G)$ under the norm $\|\cdot\|_Q$. Hence our space $Q(G)$ is isomorphic to the predual discovered in [8].

Corollary 6.6. *The map $i_Q : L^1(G) \rightarrow Q(G)$ is an injective (completely) contractive map with dense range. Moreover, for f in $L^1(G)$ and u in $M_{cb}A(G)$, we have $\langle u, i_Q(f) \rangle = \int_G u(s)f(s)ds$.*

Proof. That i_Q is contractive was shown above. That $i_Q(L^1(G))$ is dense in $Q(G)$ follows from that $L^1(G) \otimes^h L^1(G)$ is dense in $L^1(G) \otimes^h L^1(G)$. If $u \in M_{cb}A(G)$, let $u^G(s, t) = u(st)$ so $u \in \sigma V_{\text{inv}}^\infty(G)$. Then for f in $L^1(G)$, writing $f = g*h$, we have that

$$\begin{aligned} \langle u, i_Q(f) \rangle &= \langle u^G, g \otimes h \rangle = \int_G \int_G u(st)g(s)h(t)dtds \\ &= \int_G \int_G u(t)g(s)h(s^{-1}t)dtds = \int_G u(t)g*h(t)dt = \int_G u(t)f(t)dt. \end{aligned}$$

That i_Q is injective follows from that $A(G)$ is both contained in $M_{cb}A(G)$ and dense in $\mathcal{C}_0(G)$, so for every f in $L^1(G)$ there is u in $A(G)$ such that $\int_G f(t)u(t)dt \neq 0$. $\square$

We now want to see how a continuous unitary representation of G induces a map on $Q(G)$. There is an alternative approach in [29]. The author is grateful to J. Kraus for showing him this paper.

If π is a continuous unitary representation of G , let π_1 be its extension to $L^1(G)$. We have that $\pi_1 \otimes \pi_1 : L^1(G) \otimes^h L^1(G) \rightarrow C_\pi^* \otimes^h C_\pi^*$ is a complete contraction. Let $m_h : C_\pi^* \otimes^h C_\pi^* \rightarrow C_\pi^*$ be the completely contractive multiplication map. Then let

$$(6.4) \quad m_\pi = m_h \circ (\pi_1 \otimes \pi_1) : L^1(G) \otimes^h L^1(G) \rightarrow C_\pi^*.$$

Since m_π is continuous and $\pi_1 \circ m_0 = m_\pi$ on $L^1(G) \otimes^h L^1(G)$, $\ker m_\pi \supset \overline{\ker m_0}$. Hence m_π induces a map $\pi_Q : Q(G) = L^1(G) \otimes^h L^1(G) / \overline{\ker m_0} \rightarrow C_\pi^*$ by

$$(6.5) \quad \pi_Q(m(x)) = m_\pi(x)$$

for x in $L^1(G) \otimes^h L^1(G)$, where $m : L^1(G) \otimes^h L^1(G) \rightarrow Q(G)$ is the quotient map. Note that $\pi_Q^* : B_\pi \rightarrow M_{cb}A(G)$ is the completely contractive inclusion

map, so π_Q is completely contractive. Finally, note that the diagram

$$(6.6) \quad \begin{array}{ccc} L^1(G) \otimes^h L^1(G) & \xrightarrow{m_\pi} & C_\pi^* \\ & \searrow m \quad \nearrow \pi_Q & \\ & Q(G) & \end{array}$$

commutes.

It is natural to ask when π_Q is injective, since it is tempting to want to think of $Q(G)$ as a dense subspace of $C^*(G)$. However, there is a technical obstruction against obtaining this. Recall that ϖ denotes the universal representation of G , and $B(G) = A_\varpi$.

Proposition 6.7. *The map π_Q is injective if and only if A_π is weak* dense in $M_{cb}A(G)$. In particular, $\varpi_Q : Q(G) \rightarrow C^*(G)$ is injective if and only if $B(G)$ is weak* dense in $M_{cb}A(G)$.*

Proof. π_Q is injective if and only if $(C_\pi^*)^* \cong B_\pi$ is point separating for $Q(G)$. By the Hahn-Banach Theorem, this happens exactly when B_π is weak* dense in $M_{cb}A(G)$. Since A_π is $\sigma(B_\pi, C_\pi^*)$ -dense in B_π , A_π is $\sigma(B_\pi, \pi_Q(Q(G)))$ -dense in B_π , and thus its weak* closure coincides with that of B_π in $M_{cb}A(G)$. $\square$

We say that G has the *approximation property* [23] if $A(G)$ is weak* dense in $M_{cb}A(G)$. Recall that λ denotes the left regular representation of G , and $A(G) = A_\lambda$.

Corollary 6.8. *λ_Q is injective if and only if G satisfies the approximation property.*

It is natural to ask when the maps m_π and π_Q are surjective.

Proposition 6.9. *For a continuous unitary representation π of a locally compact group G , the following are equivalent.*

- (i) *The map $m_\pi : L^1(G) \otimes^h L^1(G) \rightarrow C_\pi^*$ is surjective.*
- (ii) *The map $\pi_Q : Q(G) \rightarrow C_\pi^*$ is surjective.*
- (iii) *The injection $B_\pi \hookrightarrow M_{cb}A(G)$ is bounded below.*

Proof. That (i) is equivalent to (ii) follows from (6.6). That (ii) is equivalent to (iii) follows from Theorem 6.5 and the Open Mapping Principle. $\square$

In what follows we obtain an interesting characterisation of the amenability of G which is essentially a reformulation of the fact that

$$(6.7) \quad \begin{array}{l} \text{if the injection } A(G) \hookrightarrow M_{cb}A(G) \\ \text{is bounded below, then } G \text{ is amenable.} \end{array}$$

This highly non-trivial fact was proved by V. Losert, likely in a similar vein as [33], though it remains unpublished. The author is grateful to Z.-J. Ruan for showing him this. It follows then that $B(G) = M_{cb}A(G)$ only if G is amenable. This fact is known for discrete groups by the attractive paper

[6]. In [33] it was shown that $\text{MA}(G) = \text{B}(G)$ implies that G is amenable. If G has the approximation property, then a nice proof that $\text{B}(G) = \text{M}_{cb}\text{A}(G)$ implies that G is amenable can be found in [31].

The characterisation below can also be found in [40] in the case that G is discrete.

Theorem 6.10. *The following are equivalent for any locally compact group G :*

- (i) G is amenable.
- (ii) The map $m_\varpi : L^1(G) \otimes^h L^1(G) \rightarrow C^*(G)$ is surjective.
- (iii) The map $m_\lambda : L^1(G) \otimes^h L^1(G) \rightarrow C_r^*(G)$ is surjective.

Proof. (i) $\Rightarrow$ (ii) If G is amenable, then $\text{B}(G) = \text{M}_{cb}\text{A}(G)$ isometrically. In particular, condition (iii) of Proposition 6.9 is met.

The implication (ii) $\Rightarrow$ (iii) follows from the commutativity of the following diagram.

$$\begin{array}{ccccc}
 L^1(G) \otimes^h L^1(G) & \xrightarrow{\varpi_1 \otimes \varpi_1} & C^*(G) \otimes^h C^*(G) & \xrightarrow{m_h} & C^*(G) \\
 & \searrow \lambda_1 \otimes \lambda_1 & & & \downarrow \lambda_* \\
 & & C_r^*(G) \otimes^h C_r^*(G) & \xrightarrow{m_h} & C_r^*(G)
 \end{array}$$

That (iii) $\Rightarrow$ (i) follows from Proposition 6.9 and (6.7). $\square$

We note that since $L^1(G)$ has a bounded approximate identity, the condition (ii) above is equivalent to saying that $L^1(G)$ has *similarity degree 2*, in the terminology of [40]. Furthermore, this condition implies the following result, obtained in the discrete case by Pisier in [40], and announced by him for the case of non-discrete groups in [39]:

Theorem 6.11. *A locally compact group G is amenable if and only if for every bounded representation $\pi : G \rightarrow \mathcal{B}(\mathcal{H})_{\text{inv}}$ (i.e. for which $\|\pi\|_\infty = \sup\{\|\pi(s)\| : s \in G\} < +\infty$), there is an operator S in $\mathcal{B}(\mathcal{H})_{\text{inv}}$ such that $S\pi(s)S^{-1}$ is unitary for each s in G , and $\|S\| \|S^{-1}\| \leq \|\pi\|_\infty^2$.*

The necessity condition above is from [11]. Details of Theorem 6.11 are verified in [44].

REFERENCES

- [1] G. Arsac. Sur l'espace de Banach engendr   par les coefficients d'une repr  sentation unitaire. *Pub. D  p. Math. Lyon*, 13(2):1–101, 1976.
- [2] W. Arveson. Operator algebras and invariant subspaces. *Annals of Math.*, 100:433–532, 1974.
- [3] D. P. Blecher. The standard dual of an operator space. *Pacific Math. J.*, 153(1):15–30, 1992.
- [4] D. P. Blecher and V. I. Paulsen. Tensor products of operator spaces. *J. Funct. Anal.*, 99:262–292, 1991.
- [5] D. P. Blecher and R. R. Smith. The dual of the Haagerup tensor product. *J. London Math. Soc.*, 45(2):126–144, 1992.

- [6] M. Bożejko. Positive definite bounded matrices and a characterization of amenable groups. *Proc. Amer. Math. Soc.*, 95(3):357–360, 1985.
- [7] M. Bożejko and G. Fendler. Herz-Schur multipliers and completely bounded multipliers of the Fourier algebra of a locally compact group. *Boll. Un. Mat. Ital. A (6)*, 3(2):297–302, 1984.
- [8] J. de Canniere and U. Haagerup. Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups. *Amer. J. Math.*, 107:455–500, 1984.
- [9] A. Defant and K. Floret. *Tensor Norms and Operator Ideals*, volume 176 of *Math. Studies*. North-Holland, Amsterdam, 1993.
- [10] J. Diestel and J. J. Uhl. *Vector Measures*, volume 15 of *Mathematical Surveys and Monographs*. Amer. Math. Soc., Providence, R. I., 1977.
- [11] J. Dixmier. Les moyennes invariante dans les semi-groupes et leurs applications. *Acta Sci. Math. Szeged*, 12:213–227, 1950.
- [12] J. Dixmier. *C*-algebras*, volume 15 of *North-Holland Math. Library*. North-Holland Publishing Co, 1977. Translated from the French by Francis Jellett.
- [13] E. G. Effros, J. Kraus, and Z.-J. Ruan. On two quantized tensor products. In *Operator algebras, mathematical physics, and low-dimensional topology (Istanbul, 1991)*, pages 125–145, Wellesley, MA, 1993. A K Peters.
- [14] E. G. Effros and Z.-J. Ruan. Operator convolution algebras: an approach to quantum groups. Accepted in *J. Operator Theory*, 2002.
- [15] E. G. Effros and Z.-J. Ruan. Self duality for the Haagerup tensor product and Hilbert space factorizations. *J. Funct. Anal.*, 100:257–284, 1991.
- [16] E. G. Effros and Z.-J. Ruan. *Operator Spaces*, volume 23 of *London Math. Soc., New Series*. Clarendon Press, Oxford Univ. Press, New York, 2000.
- [17] P. Eymard. L’algebre de Fourier d’un groupe localement compact. *Bull. Soc. Math. France*, 92:181–236, 1964.
- [18] J. Froelich. Compact operators, invariant subspaces and spectral synthesis. *J. Funct. Anal.*, 81:1–37, 1988.
- [19] F. Ghahramani. Isometric representations of $M(G)$ on $\mathcal{B}(\mathcal{H})$. *Glasgow Math. J.*, 23:119–122, 1982.
- [20] J. E. Gilbert. Convolution operators and Banach space tensor products I, II and III. Unpublished.
- [21] U. Haagerup. Decomposition of completely bounded maps on operator algebras. Unpublished, 1980.
- [22] U. Haagerup. Group C*-algebras without the completely bounded approximation property. Unpublished, 1988.
- [23] U. Haagerup and J. Kraus. Approximation properties for group C*-algebras and group von Neumann algebras. *Trans. Amer. Math. Soc.*, 344:667–699, 1994.
- [24] C. S. Herz. Remarques sur la note précédente de M. Varopoulos. *C. R. Acad. Sci. Paris*, 260(23):6001–6004, 1965.
- [25] C. S. Herz. Une généralisation de la notion transformée de Fourier-Stieltjes. *Ann. Inst. Fourier (Grenoble)*, 24(3):145–157, 1974.
- [26] E. Hewitt and K. A. Ross. *Abstract Harmonic Analysis II*, volume 152 of *Die Grundlehren der mathematischen Wissenschaften*. Springer, Berlin Heidelberg, 1970.
- [27] E. Hewitt and K. A. Ross. *Abstract Harmonic Analysis I*, volume 115 of *Grundlehren der mathematischen Wissenschaften*. Springer, New York, second edition, 1979.
- [28] P. Jolissaint. A characterization of completely bounded multipliers of the Fourier algebras. *Colloq. Math.*, 63:311–313, 1992.
- [29] J. Kraus. Correspondences and approximation properties for von Neumann algebras. Preprint.
- [30] J. Kraus and Z.-J. Ruan. Multipliers of Kac algebras. *Internat. J. Math.*, 8(2):213–248, 1996.

- [31] J. Kraus and Z.-J. Ruan. Approximation properties for Kac algebras. *Indiana Univ. Math. J.*, 48(2):469–535, 1999.
- [32] R. Larsen. *An Introduction to the Theory of Multipliers*, volume 175 of *Die Grundlehren der mathematischen Wissenschaften*. Springer, New York, 1971.
- [33] V. Losert. Properties of the Fourier algebra that are equivalent to amenability. *Proc. Amer. Math. Soc.*, 92(3):347–353, 1984.
- [34] M. Neufang. Isometric representations of convolution algebras as completely bounded module homomorphisms and a characterization of the measure algebra. Unpublished, 2001.
- [35] M. Neufang and N. Spronk. Completely isometric representations of measure algebras and of completely bounded multipliers of Fourier algebras. In preparation, 2003.
- [36] V. I. Paulsen. Completely bounded homomorphisms of operator algebras. *Proc. Amer. Math. Soc.*, 92:225–228, 1984.
- [37] G. Pisier. Multipliers and lacunary sets in non-amenable groups. *Amer. J. Math.*, 117:337–376, 1995.
- [38] G. Pisier. *Similarity Problems and Completely Bounded Maps*, volume 1618 of *Lec. Notes in Math.* Springer, Berlin Heidelberg, 1996.
- [39] G. Pisier. Are unitarizable groups amenable? Based on lecture at Ponza, 1999.
- [40] G. Pisier. The similarity degree of an operator algebra. *St. Petersburg Math. J.*, 10(1):103–146, 1999.
- [41] V. Runde. *Lectures on Amenability*, volume 1774 of *Lec. Notes in Math.* Springer, Berlin Heidelberg, 2002.
- [42] V. S. Shulman and L. Turowska. Operator synthesis. Preprint 2002:42, Dept. of Math., Chalmers University in Göteborg, Sweden, 2002.
- [43] R. R. Smith. Completely bounded module maps and the Haagerup tensor product. *J. Funct. Anal.*, 102(1):156–175, 1991.
- [44] N. Spronk. *On Multipliers of the Fourier Algebra of a Locally Compact Group*. PhD thesis, University of Waterloo, Waterloo, Ontario, Canada, 2002.
- [45] N. Spronk and L. Turowska. Spectral synthesis and operator synthesis for compact groups. *J. London Math. Soc.*, 66:361–376, 2002.
- [46] W.F. Stinespring. Positive functions on C^* -algebras. *Proc. Amer. Math. Soc.*, 6:211–216, 1955.
- [47] E. Størmer. Regular Abelian Banach algebras of linear maps of operator algebras. *J. Funct. Anal.*, 37:331–373, 1980.
- [48] J. Tomiyama. Tensor products of commutative Banach algebras. *Tôhoku Math. J.*, 12(2):143–154, 1960.
- [49] S. Vaes. *Locally compact quantum groups*. PhD thesis, Katholieke U. Leuven, Belgium, 2001.
- [50] N. Th. Varopoulos. Tensor algebras and harmonic analysis. *Acta. Math.*, 119:51–112, 1967.
- [51] M. Walter. Dual algebras. *Math. Scand.*, 58:77–101, 1986.
- [52] M. Walter. On a new method for defining the norm of Fourier-Stieltjes algebras. *Pacific J. Math.*, 137(1):209–223, 1989.
- [53] G. Xu. *Multipliers of the Algebras $A_p(G)$* . PhD thesis, SUNY Buffalo, NY, 1995.

DEPARTMENT OF MATHEMATICS, TEXAS A& M UNIVERSITY, COLLEGE STATION, TEXAS 77843-3368, U.S.A.

E-mail address: spronk@math.tamu.edu